

Short Review

RIFM fragrance ingredient safety assessment, isobutyl acetate, CAS Registry Number 110-19-0

A.M. Api^a, D. Belsito^b, D. Botelho^a, M. Bruze^c, G.A. Burton Jr.^d, J. Buschmann^e, M.L. Dagli^f, M. Date^a, W. Dekant^g, C. Deodhar^a, M. Francis^a, A.D. Fryer^h, L. Jones^a, K. Joshi^a, S. La Cava^a, A. Lapczynski^a, D.C. Lieblerⁱ, D. O'Brien^a, A. Patel^a, T.M. Penning^j, G. Ritacco^a, J. Romine^a, N. Sadekar^a, D. Salvito^a, T.W. Schultz^k, I.G. Sipes^l, G. Sullivan^{a,*}, Y. Thakkar^a, Y. Tokura^m, S. Tsang^a

^a Research Institute for Fragrance Materials, Inc., 50 Tice Boulevard, Woodcliff Lake, NJ, 07677, USA

^b Member RIFM Expert Panel, Columbia University Medical Center, Department of Dermatology, 161 Fort Washington Ave., New York, NY, 10032, USA

^c Member RIFM Expert Panel, Malmo University Hospital, Department of Occupational & Environmental Dermatology, Sodra Forstadsgatan 101, Entrance 47, Malmo, SE-20502, Sweden

^d Member RIFM Expert Panel, School of Natural Resources & Environment, University of Michigan, Dana Building G110, 440 Church St., Ann Arbor, MI, 48109, USA

^e Member RIFM Expert Panel, Fraunhofer Institute for Toxicology and Experimental Medicine, Nikolai-Fuchs-Strasse 1, 30625, Hannover, Germany

^f Member RIFM Expert Panel, University of Sao Paulo, School of Veterinary Medicine and Animal Science, Department of Pathology, Av. Prof. dr. Orlando Marques de Paiva, 87, Sao Paulo, CEP 05508-900, Brazil

^g Member RIFM Expert Panel, University of Wuerzburg, Department of Toxicology, Versbacher Str. 9, 97078, Wuerzburg, Germany

^h Member RIFM Expert Panel, Oregon Health Science University, 3181 SW Sam Jackson Park Rd., Portland, OR, 97239, USA

ⁱ Member RIFM Expert Panel, Vanderbilt University School of Medicine, Department of Biochemistry, Center in Molecular Toxicology, 638 Robinson Research Building, 2200 Pierce Avenue, Nashville, TN, 37232-0146, USA

^j Member of RIFM Expert Panel, University of Pennsylvania, Perelman School of Medicine, Center of Excellence in Environmental Toxicology, 1316 Biomedical Research Building (BRB) II/III, 421 Curie Boulevard, Philadelphia, PA, 19104-3083, USA

^k Member RIFM Expert Panel, The University of Tennessee, College of Veterinary Medicine, Department of Comparative Medicine, 2407 River Dr., Knoxville, TN, 37996-4500, USA


^l Member RIFM Expert Panel, Department of Pharmacology, University of Arizona, College of Medicine, 1501 North Campbell Avenue, P.O. Box 245050, Tucson, AZ, 85724-5050, USA

^m Member RIFM Expert Panel, The Journal of Dermatological Science (JDS), Editor-in-Chief, Professor and Chairman, Department of Dermatology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, 431-3192, Japan

Version: 073018. This version replaces any previous versions.

Name: Isobutyl acetate

CAS Registry Number: 110-19-0

**Abbreviation/Definition List:**

2-Box Model - A RIFM, Inc. proprietary *in silico* tool used to calculate fragrance air exposure concentration

AF - Assessment Factor

BCF - Bioconcentration Factor

Crema RIFM Model - The Crema RIFM Model uses probabilistic (Monte Carlo) simulations to allow full distributions of data sets, providing a more realistic estimate of aggregate exposure to individuals across a population (Comiskey et al., 2015, 2017; Safford et al., 2015, 2017) compared to a deterministic aggregate approach

DEREK - Derek Nexus is an *in silico* tool used to identify structural alerts

DST - Dermal Sensitization Threshold

ECHA - European Chemicals Agency

EU - Europe/European Union

GLP - Good Laboratory Practice

IFRA - The International Fragrance Association

LOEL - Lowest Observable Effect Level

* Corresponding author.

E-mail address: gsullivan@rifm.org (G. Sullivan).

<https://doi.org/10.1016/j.fct.2019.04.058>

Received 23 August 2018; Received in revised form 6 March 2019; Accepted 30 April 2019

Available online 06 May 2019

0278-6915/ © 2019 Elsevier Ltd. All rights reserved.

MOE - Margin of Exposure
MPPD - Multiple-Path Particle Dosimetry. An *in silico* model for inhaled vapors used to simulate fragrance lung deposition
NA - North America
NESIL - No Expected Sensitization Induction Level
NOAEC - No Observed Adverse Effect Concentration
NOAEL - No Observed Adverse Effect Level
NOEC - No Observed Effect Concentration
NOEL - No Observed Effect Level
OECD - Organisation for Economic Co-operation and Development
OECD TG - Organisation for Economic Co-operation and Development Testing Guidelines
PBT - Persistent, Bioaccumulative, and Toxic
PEC/PNEC - Predicted Environmental Concentration/Predicted No Effect Concentration
QRA - Quantitative Risk Assessment
REACH - Registration, Evaluation, Authorisation, and Restriction of Chemicals
RfD - Reference Dose
RIFM - Research Institute for Fragrance Materials
RQ - Risk Quotient
Statistically Significant - Statistically significant difference in reported results as compared to controls with a $p < 0.05$ using appropriate statistical test.
TTC - Threshold of Toxicological Concern
UV/Vis Spectra - Ultraviolet/Visible Spectra
VCF - Volatile Compounds in Food
VoU - Volume of Use
vPvB - (very) Persistent, (very) Bioaccumulative
WoE - Weight of Evidence

The Expert Panel for Fragrance Safety* concludes that this material is as described in this safety assessment.

This safety assessment is based on the RIFM Criteria Document (Api et al., 2015), which should be referred to for clarifications.

Each endpoint discussed in this safety assessment includes the relevant data that were available at the time of writing (version number in the top box is indicative of the date of approval based on a 2-digit month/day/year), both in the RIFM database (consisting of publicly available and proprietary data) and through publicly available information sources (e.g., SciFinder and PubMed). Studies selected for this safety assessment were based on appropriate test criteria, such as acceptable guidelines, sample size, study duration, route of exposure, relevant animal species, most relevant testing endpoints, etc. A key study for each endpoint was selected based on the most conservative endpoint value (e.g., PNEC, NOAEL, LOEL, and NESIL).

*The Expert Panel for Fragrance Safety is an independent body that selects its own members and establishes its own operating procedures. The Expert Panel is comprised of internationally known scientists that provide RIFM with guidance relevant to human health and environmental protection.

Summary: The existing information supports the use of this material as described in this safety assessment.

Isobutyl acetate (CAS # 110-19-0) was evaluated for genotoxicity, repeated dose toxicity, developmental and reproductive toxicity, local respiratory toxicity, phototoxicity/photoallergenicity, skin sensitization, and environmental safety. Data from isobutyl acetate show that it is not (expected to be) genotoxic. Data from the target material and read-across analog isoamyl acetate (CAS # 123-92-2) show that there are no safety concerns for isobutyl acetate for skin sensitization under the current declared levels of use. The repeated dose and reproductive endpoints were evaluated using the TTC for a Cramer Class I material and the exposure to isobutyl acetate is below the TTC (0.03 mg/kg/day and 0.03 mg/kg/day, respectively). For the local respiratory endpoint, a calculated MOE > 100 was provided by the read-across analog butyl acetate (CAS # 123-86-4). The phototoxicity/photoallergenicity endpoints were evaluated based on UV spectra; isobutyl acetate is not expected to be phototoxic/photoallergenic. For the environmental endpoints, isobutyl acetate is not a PBT as per the IFRA Environmental Standards, and its risk quotients (i.e., PEC/PNEC) for the aquatic environment based on the screening-level is < 1.

Human Health Safety Assessment

Genotoxicity: Not genotoxic. (ECHA, 2011)

Repeated Dose Toxicity: No NOAEL available. Exposure is below the TTC.

Reproductive Toxicity: No NOAEL available. Exposure is below the TTC.

Skin Sensitization: No safety concerns at current, declared use levels. (RIFM, 1987)

Phototoxicity/Photoallergenicity: Not expected to be phototoxic/photoallergenic. (UV Spectra, RIFM Database)

Local Respiratory Toxicity: NOAEC = 2375 mg/m³.

(ECHA, 2011; accessed 11/30/17; data also available in David et al., 2001)

Environmental Safety Assessment

Hazard Assessment:

Persistence: Critical Measured Value: 81% (OECD 301C) (EPI Suite v4.1; US EPA, 2012a)

Bioaccumulation: Screening-level: 6.49 L/kg (EPI Suite v4.1; US EPA, 2012a)

Ecotoxicity: Screening-level: Fish LC50: 258.6 mg/L (RIFM Framework; Salvito et al., 2002)

Conclusion: Not PBT or vPvB as per IFRA Environmental Standards

Risk Assessment:

Screening-level: PEC/PNEC (North America and Europe) < 1 (RIFM Framework; Salvito et al., 2002)

Critical Ecotoxicity Endpoint: Fish LC50: 258.6 mg/L (RIFM Framework; Salvito et al., 2002)

RIFM PNEC is: 0.2856 µg/L

● **Revised PEC/PNECs (2015 IFRA VoU):** North America and Europe: Not applicable; cleared at the screening-level

1. Identification

- 1. Chemical Name:** Isobutyl acetate
- 2. CAS Registry Number:** 110-19-0
- 3. Synonyms:** Acetic acid, 2-methylpropyl ester; Isobutyl ethanoate; 2-Methyl-1-propyl acetate; 2-Methylpropyl acetate; Acetic acid, isobutyl ester; 酢酸ブチル; Isobutyl acetate

4. Molecular Formula: C₆H₁₂O₂

5. Molecular Weight: 116.16

6. RIFM Number: 811

7. Stereochemistry: Isomer not specified. No stereocenter and no stereoisomers possible.

2. Physical data

- Boiling Point:** 117 °C (FMA Database), 111.74 °C (EPI Suite)
- Flash Point:** 22 °C (GHS), 65 °F; CC (FMA Database)
- Log K_{ow}:** 1.60 (Abraham and Rafols, 1995), 1.77 (EPI Suite), partition coefficient in water/air = 25.9 (SD 2.1) (Kaneko et al., 1994)
- Melting Point:** −68.43 °C (EPI Suite)
- Water Solubility:** 3128 mg/L (EPI Suite)
- Specific Gravity:** 0.8676 (EOA Sample 76–164), 0.87 (FMA)
- Vapor Pressure:** 13.7 mm Hg @ 20 °C (EPI Suite), 13 mm Hg @ 20 °C (FMA Database), 18.3 mm Hg @ 25 °C (EPI Suite)
- UV Spectra:** No significant absorbance between 290 and 700 nm; molar absorption coefficient is below the benchmark (1000 L mol^{−1} · cm^{−1})
- Appearance/Organoleptic:** A colorless liquid with a sweet, tropical, and fruity odor*

*<http://www.thegoodscentscompany.com/data/rw1013631.html>, 07/30/18.

3. Exposure

- Volume of Use (worldwide band):** 1–100 metric tons per year (IFRA, 2015)
- 95th Percentile Concentration in Hydroalcohols:** 0.024% (RIFM, 2017)
- Inhalation Exposure*:** 0.00023 mg/kg/day or 0.017 mg/day (RIFM, 2017)
- Total Systemic Exposure**:** 0.0024 mg/kg/day (RIFM, 2017)

*95th percentile calculated exposure derived from concentration survey data in the Creme RIFM aggregate exposure model (Comiskey et al., 2015, 2017; Safford et al., 2015, 2017).

**95th percentile calculated exposure; assumes 100% absorption unless modified by dermal absorption data as reported in Section IV. It is derived from concentration survey data in the Creme RIFM aggregate exposure model and includes exposure via dermal, oral, and inhalation routes whenever the fragrance ingredient is used in products that include these routes of exposure (Comiskey et al., 2015, 2017; Safford et al., 2015, 2017).

4. Derivation of systemic absorption

- Dermal:** Assumed 100%
- Oral:** Assumed 100%
- Inhalation:** Assumed 100%

5. Computational toxicology evaluation

- Cramer Classification:** Class I, Low

Expert Judgment	Toxtree v 2.6	OECD QSAR Toolbox v 3.2
I	I	I

2. Analogs Selected:

- Genotoxicity:** None
 - Repeated Dose Toxicity:** None
 - Reproductive Toxicity:** None
 - Skin Sensitization:** Isoamyl acetate (CAS # 123-92-2)
 - Phototoxicity/Photoallergenicity:** None
 - Local Respiratory Toxicity:** Butyl acetate (CAS # 123-86-4)
 - Environmental Toxicity:** None
- Read-across Justification:** See Appendix below

6. Metabolism

No relevant data available for inclusion in this safety assessment.

7. Natural occurrence (discrete chemical) or composition (NCS)

Isobutyl acetate is reported to occur in nature in the following*:

Acerola (*Malpighia*)
 Alpinia species
 Apple brandy (*Calvados*)
 Apple fresh (*Malus* species)
 Apple processed (*Malus* species)
 Apricot (*Prunus armeniaca* L.)
 Arrack
 Banana (*Musa sapientum* L.)
 Bantu beer
 Beef
 Beer
 Black currants (*Ribes nigrum* L.)
 Camomile
 Cape gooseberry (*Physalis peruviana* L.)
 Cashew apple wine
 Ceriman, pinanona (*Monstera deliciosa* Liebm.)
 Cheese, various types
 Cider (apple wine)
 Cocoa category
 Fig (*Ficus carica* L.)
 Grape (*Vitis* species)
 Grape brandy
 Guava and feyoa
 Guava wine
 Honey
 Laurel (*Laurus nobilis* L.)
 Litchi (*Litchi chinensis* Sonn.)
 Litchi wine
Mangifera species
 Melon
 Miso (soy bean, rice, or fish)
 Naranjilla fruit (*Solanum quitoense* Lam.)
 Nectarine
 Olive (*Olea europaea*)
 Papaya (*Carica papaya* L.)
 Passion fruit (*Passiflora* species)
 Peach (*Prunus persica* L.)
 Pear (*Pyrus communis* L.)
 Pear brandy
 Pepino fruit (*Solanum muricatum*)
 Pineapple (*Ananas comosus*)
 Pistachio nut (*Pistacia vera*)
 Plum (*Prunus* species)
 Plum brandy
 Plum wine
 Pomegranate wine (*Punica granatum* L.)
 Quince, marmelo (*Cydonia oblonga* Mill.)
 Rambutan (*Nephelium lappaceum* L.)
 Rum
 Sake
 Sherry
 Shoyu (fermented soya hydrolysate)
 Spineless monkey orange (*Strychnos madagasc.*)
 Starfruit (*Averrhoa carambola* L.)
 Strawberry (*Fragaria* species)
 Tamarind (*Tamarindus indica* L.)
 Tomato (*Lycopersicon esculentum* Mill.)
 Vinegar
 Wheat bread

Whisky

Wine

*VCF Volatile Compounds in Food: Database/Nijssen, L.M.; Ingen-Visscher, C.A. van; Donders, J.J.H. (eds). – Version 15.1 – Zeist (The Netherlands): TNO Triskelion, 1963–2014. A continually updated database containing information on published volatile compounds that have been found in natural (processed) food products. Includes FEMA GRAS and EU-Flavis data.

8. IFRA standard

None

9. REACH dossier

Available, accessed 07/31/18 (ECHA, 2011)

10. Summary

10.1. Human health endpoint summaries

10.1.1. Genotoxicity

Based on the current existing data and use levels, isobutyl acetate does not present a concern for genetic toxicity.

10.1.1.1. Risk assessment. Isobutyl acetate was assessed in the BlueScreen assay and found negative for both cytotoxicity (positive: < 80% relative cell density) and genotoxicity, with and without metabolic activation (RIFM, 2013). BlueScreen is a screening assay which assesses genotoxic stress through human derived gene expression. Additional assays were considered to fully assess the potential mutagenic or clastogenic effects on the target material.

The mutagenic activity of isobutyl acetate has been evaluated in a bacterial reverse mutation assay conducted in compliance with GLP regulations and according to guidelines similar to OECD TG 471 using the standard plate incorporation method. *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and TA1538 were treated with isobutyl acetate in dimethyl sulfoxide (DMSO) at concentrations up to 10000 µg/plate. No increases in the mean number of revertant colonies were observed at any tested dose in the presence or absence of S9 (ECHA, 2011). Under the conditions of the study, isobutyl acetate was not mutagenic in the Ames test.

The clastogenicity of isobutyl acetate was assessed in an *in vitro* chromosome aberration study conducted in compliance with GLP regulations and in accordance with OECD TG 473. Chinese hamster lung fibroblasts (V79 cells) were treated with isobutyl acetate in MEM culture medium at concentrations up to 5000 µg/mL in the presence and absence of exogenous metabolic activation. No significant increases in the frequency of cells with structural chromosomal aberrations or polyploid cells were observed with any dose of the test item, either with or without S9 metabolic activation (ECHA, 2011). Under the conditions of the study, isobutyl acetate was considered to be non-clastogenic in the *in vitro* chromosome aberration assay.

Based on the data available, isobutyl acetate does not present a concern for genotoxic potential.

Additional References: OECD SIDS Database

Literature Search and Risk Assessment Completed On: 12/5/2017

10.1.2. Repeated dose toxicity

There are insufficient repeated dose toxicity data on isobutyl acetate or any read-across materials. The total systemic exposure to isobutyl acetate is below the TTC for the repeated dose toxicity endpoint of a Cramer Class I material at the current level of use.

10.1.2.1. Risk assessment. There are no repeated dose toxicity data on

isobutyl acetate or any read-across materials that can be used to support the repeated dose toxicity endpoint. The total systemic exposure to isobutyl acetate (2.4 µg/kg/day) is below the TTC (30 µg/kg bw/day; Kroes et al., 2007) for the repeated dose toxicity endpoint of a Cramer Class I material at the current level of use.

Additional References: None

Literature Search and Risk Assessment Completed On: 11/21/17

10.1.3. Reproductive toxicity

There are insufficient reproductive toxicity data on isobutyl acetate or any read-across materials. The total systemic exposure to isobutyl acetate is below the TTC for the reproductive toxicity endpoint of a Cramer Class I material at the current level of use.

10.1.3.1. Risk assessment. There are no reproductive toxicity data on isobutyl acetate or any read-across materials that can be used to support the reproductive toxicity endpoint. The total systemic exposure to isobutyl acetate (2.4 µg/kg bw/day) is below the TTC (30 µg/kg bw/day; Kroes et al., 2007; Laufersweiler et al., 2012) for the reproductive toxicity endpoint of a Cramer Class I material at the current level of use.

Additional References: None

Literature Search and Risk Assessment Completed On: 11/21/17

10.1.4. Skin sensitization

Based on the existing data and read-across isoamyl acetate (CAS # 123-92-2), isobutyl acetate does not present a safety concern for skin sensitization under the current, declared levels of use.

10.1.4.1. Risk assessment. Limited skin sensitization studies are available for isobutyl acetate. Based on the existing data and read-across isoamyl acetate (CAS # 123-92-2; see Section V), isobutyl acetate does not present a safety concern for skin sensitization under the current, declared levels of use. The chemical structure of these materials indicate that they would not be expected to react with skin proteins (Toxtree 2.6.13; OECD toolbox v3.4). In guinea pigs, maximization tests and an open epicutaneous test with read-across isoamyl acetate did not present reactions indicative of sensitization (Ballantyne et al., 1986; Klecak, 1985). In a human maximization test, no skin sensitization reactions were observed with isobutyl acetate and read-across isoamyl acetate (RIFM, 1976; RIFM, 1973). Additionally, in a confirmatory human repeat insult patch test (HR IPT) with 23622 µg/cm² of read-across material isoamyl acetate, no reactions indicative of sensitization were observed in any of the 197 volunteers (RIFM, 1987).

Based on weight of evidence from structural analysis, animal and human studies, and read-across isoamyl acetate, isobutyl acetate does not present a safety concern for skin sensitization under the current, declared levels of use.

Additional References: None

Literature Search and Risk Assessment Completed On: 11/13/17

10.1.5. Phototoxicity/photoallergenicity

Based on UV/Vis absorption spectra, isobutyl acetate would not be expected to present a concern for phototoxicity or photoallergenicity.

10.1.5.1. Risk assessment. There are no phototoxicity studies available for isobutyl acetate in experimental models. UV/Vis absorption spectra indicate no significant absorption between 290 and 700 nm. The corresponding molar absorption coefficient is well below the benchmark of concern for phototoxicity and photoallergenicity (Henry et al., 2009). Based on lack of absorbance, isobutyl acetate does not present a concern for phototoxicity or photoallergenicity.

10.1.5.2. UV spectra analysis. UV/Vis absorption spectra (OECD TG

101) were obtained. The spectra indicate no significant absorbance in the range of 290–700 nm. The molar absorption coefficient is below the benchmark of concern for phototoxic effects, $1000 \text{ L mol}^{-1} \cdot \text{cm}^{-1}$ (Henry et al., 2009).

Additional References: None

Literature Search and Risk Assessment Completed On: 11/30/17

10.1.6. Local respiratory toxicity

There is insufficient inhalation data available on isobutyl acetate; however, in a 13-week inhalation study for the read-across analog butyl acetate (CAS # 123-86-4; see section V), a NOAEC of 2375 mg/m^3 is reported (ECHA, 2011; accessed 08/03/17; David et al., 2001).

10.1.6.1. Risk assessment. The inhalation exposure estimated for combined exposure was considered along with toxicological data observed in the scientific literature to calculate the MOE for local respiratory toxicity. In a 13-week whole-body inhalation study conducted in rats, a NOAEC of 2375 mg/m^3 (500 ppm) was reported (ECHA, 2011; accessed 11/30/17; David et al., 2001). Whole-body inhalation exposure of read-across material butyl acetate was administered at target concentrations (0 [sham], 2375, 7126, 14253 mg/m^3) to both male and female Sprague Dawley rats (15 animals/sex/concentration). Clinical observations, body weight, food consumption, ophthalmology, hematology, clinical chemistry, organ weights, gross pathology, and histopathology were all considered. Body weights and food consumption decreased among animals in mid- and high-concentration treatment groups. Organ weight changes were also dependent upon treatment and concentration. Lung weights increased among males exposed to 14253 mg/m^3 butyl acetate compared to the control group. Additionally, histopathology for both the mid- and high-concentration treatment groups demonstrated degenerated olfactory epithelial tissue as well as dorsal medial meatus and ethmotubines of the nasal passages. Severity of the histopathological findings ranged from mild to moderate for the high-concentration group, but minimal to mild for the mid-concentration group. As there were no observable adverse effects documented for the low-concentration treatment group, the NOAEC was determined to be 2375 mg/m^3 .

This NOAEC expressed in $\text{mg/kg lung weight/day}$ is:

- $(2375 \text{ mg/m}^3) (1\text{m}^3/1000 \text{ L}) = 2.375 \text{ mg/L}$
- Minute ventilation (MV) of 0.17 L/min for a Sprague Dawley rat x duration of exposure of 360 min per day (min/day) (according to GLP study guidelines) = 61.2 L/day
- $(2.375 \text{ mg/L}) (61.2 \text{ L/day}) = 145.35 \text{ mg/day}$
- $(145.35 \text{ mg/day}) / (0.0016 \text{ kg lung weight of rat}^*) = 90844 \text{ mg/kg lung weight/day}$

The 95th percentile calculated exposure to isobutyl acetate was reported to be 0.017 mg/day —this value was derived from the concentration survey data in the Creme RIFM Exposure Model (Comiskey et al., 2015; Safford et al., 2015). To compare this estimated exposure with the NOAEC expressed in $\text{mg/kg lung weight/day}$, this value is divided by $0.65 \text{ kg human lung weight}$ (Carthew et al., 2009) to give $0.026 \text{ mg/kg lung weight/day}$ resulting in an MOE of 3494000 (i.e., $[90844 \text{ mg/kg lung weight/day}] / [0.026 \text{ mg/kg lung weight/day}]$).

The MOE is greater than 100. Without adjustment for specific uncertainty factors related to inter-species and intra-species variation, the material exposure by inhalation at 0.017 mg/day is deemed to be safe under the most conservative consumer exposure scenario.

* Phalen, R.F. Inhalation Studies. Foundations and Techniques, 2nd Ed 2009. Published by, Informa Healthcare USA, Inc., New York,

NY. Chapter 9, Animal Models, in section: “Comparative Physiology and Anatomy,” subsection, “Comparative Airway Anatomy.”

Additional References: Smyth et al. (1962); Martinez-Alfaro et al. (2009); Smyth et al. (1954); Smyth and Smyth (1928); Haglund et al. (1980); Nelson et al. (1943); McOmie and Anderson. (1949); NIOSH, 1982; Burleigh-Flayer et al. (1991); Querci and Mascia. (1970); Ambrosio and D'Arrigo (1962); Ambrosio et al. (1962); Frantik et al. (1994); Querci et al. (1970); Osina (1959); Sayers et al. (1936); Iregren et al. (1993); Ashley and Prah. (1997); Bowen and Balster. (1997); Norris et al. (1997); Silver (1992); Prah et al. (1998); David et al. (1998); Kodak (1996); UnionCarbide (1993); Saillenfait et al., (2007).

Literature Search and Risk Assessment Completed On: 12/01/2017

10.2. Environmental endpoint summary

10.2.1. Screening-level assessment

A screening-level risk assessment of isobutyl acetate was performed following the RIFM Environmental Framework (Salvito et al., 2002), which provides 3 tiered levels of screening for aquatic risk. In Tier 1, only the material's regional VoU, its log K_{OW} , and its molecular weight are needed to estimate a conservative risk quotient (RQ), expressed as the ratio Predicted Environmental Concentration/Predicted No Effect Concentration (PEC/PNEC). A general QSAR with a high uncertainty factor applied is used to predict fish toxicity, as discussed in Salvito et al. (2002). In Tier 2, the RQ is refined by applying a lower uncertainty factor to the PNEC using the ECOSAR model (US EPA, 2012b), which provides chemical class-specific ecotoxicity estimates. Finally, if necessary, Tier 3 is conducted using measured biodegradation and ecotoxicity data to refine the RQ, thus allowing for lower PNEC uncertainty factors. The data for calculating the PEC and PNEC for this safety assessment are provided in the table below. For the PEC, the range from the most recent IFRA Volume of Use Survey is reviewed. The PEC is then calculated using the actual regional tonnage, not the extremes of the range. Following the RIFM Environmental Framework, isobutyl acetate was identified as a fragrance material with no potential to present a possible risk to the aquatic environment (i.e., its screening-level PEC/PNEC < 1).

A screening-level hazard assessment using EPI Suite v4.1 (US EPA, 2012a) did not identify Isobutyl acetate as either persistent or bioaccumulative based on its structure and physical-chemical properties. This screening-level hazard assessment considers the potential for a material to be persistent and bioaccumulative and toxic, or very persistent and very bioaccumulative as defined in the Criteria Document (Api et al., 2015). As noted in the Criteria Document, the screening criteria applied are the same as those used in the EU for REACH (ECHA, 2012). For persistence, if the EPI Suite model BIOWIN 3 predicts a value < 2.2 and either BIOWIN 2 or BIOWIN 6 predicts a value < 0.5, then the material is considered potentially persistent. A material would be considered potentially bioaccumulative if the EPI Suite model BCFBAF predicts a fish BCF $\geq 2000 \text{ L/kg}$. Ecotoxicity is determined in the above screening-level risk assessment. If, based on these model outputs (Step 1), additional assessment is required, a WoE-based review is then performed (Step 2). This review considers available data on the material's physical-chemical properties, environmental fate (e.g., OECD Guideline biodegradation studies or die-away studies), fish bioaccumulation, and higher-tier model outputs (e.g., US EPA's BIOWIN and BCFBAF found in EPI Suite v4.1). Data on persistence and bioaccumulation are reported below and summarized in the Environmental Safety Assessment section prior to Section 1.

10.2.2. Risk assessment

Based on the current Volume of Use (2015), isobutyl acetate does not present a risk to the aquatic compartment in the screening-level assessment.

10.2.2.1. *Biodegradation*. No data available.

10.2.2.2. *Ecotoxicity*. No data available.

10.2.2.3. *Other available data*. Isobutyl acetate has been registered under REACH and the following additional data is available:

Ready biodegradability of the test material was evaluated according to the OECD 301D method. Biodegradation of 81% was observed after 20 days.

A Fish (*Oryzias latipes*) acute toxicity study was conducted according to the OECD 203 guidelines under semi-static conditions. The 96-h LC50 was reported to be 16.6 mg/L.

A *Daphnia magna* immobilization test was conducted according to the OECD 202 method under semi-static conditions. The 48-h EC50 based on time weighted average was reported to be 24.6 mg/L.

An algae growth inhibition test was conducted according to the OECD 201 method. The 72-h EC50 was reported to be 397 mg/L and 246 mg/L based on growth rate and biomass, respectively.

A *Daphnia magna* reproduction study was conducted according to the OECD 211 method under semi-static conditions. The 21-day NOEC was reported to be 23.2 mg/L.

10.2.3. Risk assessment refinement

Since isobutyl acetate has passed the screening criteria, measured data is included for completeness only and has not been used in PNEC derivation.

Ecotoxicological data and PNEC derivation (all endpoints reported in mg/L; PNECs in µg/L)

Endpoints used to calculate PNEC are underlined.

	LC50 (Fish) (mg/L)	EC50 (<i>Daphnia</i>) (mg/L)	EC50 (Algae) (mg/L)	AF	PNEC (µg/L)	Chemical Class
RIFM Framework Screening-level (Tier 1)	<u>285.6</u>			1,000,000	0.2856	

Exposure information and PEC calculation (following RIFM Framework: [Salvito et al., 2002](#))

Exposure	Europe	North America
Log K _{ow} used	1.1	1.1
Biodegradation Factor Used	0	0
Dilution Factor	3	3
Regional Volume of Use Tonnage Band	1–10	1–10

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fct.2019.04.058>.

Risk Characterization: PEC/PNEC < 1 < 1

Based on the available data, the RQ for this material is < 1. No further assessment is necessary.

The RIFM PNEC is 0.2856 µg/L. The revised PEC/PNECs for EU and NA are: not applicable. The material was cleared at the screening-level and therefore does not present a risk to the aquatic environment at the current reported volumes of use.

Literature Search and Risk Assessment Completed On: 11/30/17

11. Literature Search*

- **RIFM Database:** Target, Fragrance Structure Activity Group materials, other references, JECFA, CIR, SIDS
- **ECHA:** <http://echa.europa.eu/>
- **NTP:** <https://ntp.niehs.nih.gov/>
- **OECD Toolbox**
- **SciFinder:** <https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf>
- **PubMed:** <http://www.ncbi.nlm.nih.gov/pubmed>
- **TOXNET:** <http://toxnet.nlm.nih.gov/>
- **IARC:** <http://monographs.iarc.fr>
- **OECD SIDS:** <http://webnet.oecd.org/hpv/ui/Default.aspx>
- **EPA ACToR:** <https://actor.epa.gov/actor/home.xhtml>
- **US EPA HPVIS:** https://ofmpub.epa.gov/opthpv/public_search_publicdetails?submission_id=24959241&ShowComments=Yes&sqlstr=null&recordcount=0&User_title=DetailQuery%20Results&EndPointRpt=Y#submission
- **Japanese NITE:** <http://www.safe.nite.go.jp/english/db.html>
- **Japan Existing Chemical Data Base (JECDB):** http://dra4.nihs.go.jp/mhlw_data/jsp/SearchPageENG.jsp
- **Google:** <https://www.google.com>
- **ChemIDplus:** <https://chem.nlm.nih.gov/chemidplus/>

Search keywords: CAS number and/or material names

*Information sources outside of RIFM's database are noted as appropriate in the safety assessment. This is not an exhaustive list. The links listed above were active as of 07/30/2018

Conflicts of interest

The authors declare that they have no conflicts of interest.




Appendix

Read-across Justification

Methods

The read-across analogs were identified following the strategy for structuring and reporting a read-across prediction of toxicity described in Schultz et al. (2015). The strategy is also consistent with the guidance provided by OECD within Integrated Approaches for Testing and Assessment (OECD, 2015) and the European Chemical Agency read-across assessment framework (ECHA, 2016).

- First, materials were clustered based on their structural similarity. Second, data availability and data quality on the selected cluster were examined. Third, appropriate read-across analogs from the cluster were confirmed by expert judgment.
- Tanimoto structure similarity scores were calculated using FCFC4 fingerprints (Rogers and Hahn, 2010).
- The physical–chemical properties of the target substance and the read-across analogs were calculated using EPI Suite (US EPA, 2012a).
- J_{\max} values were calculated using RIFM's skin absorption model (SAM). The parameters were calculated using the consensus model (Shen et al., 2014).
- DNA binding, mutagenicity, genotoxicity alerts and oncologic classification predictions were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- ER binding and repeat dose categorization were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- Developmental toxicity was predicted using CAESAR v2.1.7 (Cassano et al., 2010), and skin sensitization was predicted using Toxtree 2.6.13.
- Protein binding was predicted using OECD QSAR Toolbox v3.4 (OECD, 2012).
- The major metabolites for the target and read-across analogs were determined and evaluated using OECD QSAR Toolbox v3.4 (OECD, 2012).

	Target Material	Read-across Material	Read-across Material
Principal Name	Isobutyl acetate	Isoamyl acetate	Butyl acetate
CAS No.	110-19-0	123-92-2	123-86-4
Structure			
Similarity (Tanimoto Score)		0.64	0.75
Read-across Endpoint		<ul style="list-style-type: none"> • Skin sensitization 	<ul style="list-style-type: none"> • Local Respiratory toxicity
Molecular Formula	C ₆ H ₁₂ O ₂	C ₇ H ₁₄ O ₂	C ₆ H ₁₂ O ₂
Molecular Weight	116.16	130.19	116.16
Melting Point (°C, EPI Suite)	-68.43	-56.05	-56.83
Boiling Point (°C, EPI Suite)	111.74	134.87	125.79
Vapor Pressure (Pa @ 25°C, EPI Suite)	2.44E+003	7.47E+002	1.59E+003
Log Kow (KOWWIN v1.68 in EPI Suite)	1.78	2.25	1.78
Water Solubility (mg/L, @ 25°C, WSKOW v1.42 in EPI Suite)	6300	2000	8400
J_{\max} (µg/cm ² /h, SAM)	225.843	101.618	301.124
Henry's Law (Pa·m ³ /mol, Bond Method, EPI Suite)	4.16E+001	5.52E+001	4.16E+001
Skin Sensitization			
Protein Binding (OASIS v1.1)		<ul style="list-style-type: none"> • No alert found 	
Protein Binding (OECD)		<ul style="list-style-type: none"> • No alert found 	
Protein Binding Potency		<ul style="list-style-type: none"> • Not possible to classify 	
Protein Binding Alerts for Skin Sensitization (OASIS v1.1)	<ul style="list-style-type: none"> • No alert found 	<ul style="list-style-type: none"> • No alert found 	
Skin Sensitization Reactivity Domains (Toxtree v2.6.13)	<ul style="list-style-type: none"> • No alert found 	<ul style="list-style-type: none"> • No alert found 	
Local Respiratory Toxicity			
Respiratory Sensitization (OECD QSAR Toolbox v3.4)	<ul style="list-style-type: none"> • No alert found 		<ul style="list-style-type: none"> • No alert found
Metabolism			
Rat Liver S9 Metabolism Simulator and Structural Alerts for Metabolites (OECD QSAR Toolbox v3.4)	See Supplemental Data 1	See Supplemental Data 2	See Supplemental Data 3

Summary

There are insufficient toxicity data on isobutyl acetate (CAS # 110-19-0). Hence, *in silico* evaluation was conducted to determine read-across analogs for this material. Based on structural similarity, reactivity, metabolism, physical–chemical properties, and expert judgment, isoamyl acetate (CAS # 123-92-2) and butyl acetate (CAS # 123-86-4) were identified as read-across materials with sufficient data for toxicological evaluation.

Conclusions

- Isoamyl acetate (CAS # 123-92-2) was used as a read-across analog for the target material isobutyl acetate (CAS # 110-19-0) for the skin sensitization endpoint.
 - o The target substance and the read-across analog are structurally similar and belong to branched saturated esters.
 - o The target substance and the read-across analog share an acetate ester structure.
 - o The key structural difference between the target substance and the read-across analog is the presence of an additional methylene group in the C5 isoamyl fragment in the read-across material compared to the C4 isobutyl group in the target substance. This structural difference is toxicologically insignificant.

- o Structural similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score is mainly driven by the near identity of these branched acetate esters. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
- o The physical–chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
- o According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
- o The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
- o The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.
- Butyl acetate (CAS # 123-86-4) was used as a read-across analog for the target material isobutyl acetate (CAS # 110-19-0) for the local respiratory toxicity endpoint.
 - o The target substance and the read-across analog are structurally similar and belong to aliphatic saturated esters.
 - o The target substance and the read-across analog share an acetyl ester structure.
 - o The key structural difference between the target substance and the read-across analog is that the target substance is an ester of a branched chain C4 alcohol, whereas the read-across analog is the ester of a straight chain C4 alcohol. This structural difference is toxicologically insignificant.
 - o Structural similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score is mainly driven by the near identity of these acetate esters. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
 - o The physical–chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
 - o According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
 - o The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
 - o The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.

References

- Abraham, M.H., Rafols, C., 1995. Factors that influence tadpole narcosis. An LFER analysis. *J. Chem. Soc. Perkin Trans. 2* (10), 1843–1851.
- Ambrosio, L., D'Arrigo, S., 1962. Anatomic and pathological changes during experimental intoxication and amyl, propyl, and butyl acetates. *Folia Med.* 45, 525–537.
- Ambrosio, L., Insera, A., Bruni, D., 1962. The blood picture in amyl, butyl and propyl acetate poisoning. *Folia Med.* 45 (8), 700–717.
- Api, A.M., Belsito, D., Bruze, M., Cadby, P., Calow, P., Dagli, M.L., Dekant, W., Ellis, G., Fryer, A.D., Fukayama, M., Griem, P., Hickey, C., Kromidas, L., Lalko, J.F., Liebler, D.C., Miyachi, Y., Politano, V.T., Renskers, K., Ritacco, G., Salvito, D., Schultz, T.W., Sipes, I.G., Smith, B., Vitale, D., Wilcox, D.K., 2015. Criteria for the research institute for fragrance materials, Inc. (RIFM) safety evaluation process for fragrance ingredients. *Food Chem. Toxicol.* 82, S1–S19.
- Ashley, D.L., Prah, J.D., 1997. Time dependence of blood concentrations during and after exposure to a mixture of volatile organic compounds. *Arch. Environ. Health* 52 (1), 26–33.
- Ballantyne, B., Tyler, T.R., Auletta, C.S., 1986. The sensitizing potential of primary amyl acetate in the Guinea pig. *Vet. Hum. Toxicol.* 28 (3), 213–215.
- Bowen, S.E., Balster, R.L., 1997. A comparison of the acute behavioral effects of inhaled amyl, ethyl, and butyl acetate in mice. *Fundam. Appl. Toxicol.* 35 (2), 189–196.
- Burleigh-Player, H.D., Dodd, D.E., Walker, J.C., Jennings, R.A., Mosberg, A.T., Ogden, M.W., 1991. The respiratory effects of n-amyl and n-butyl acetate in mice. *Toxicologist* 11 (1), 86.
- Carthew, P., Clapp, C., Gutsell, S., 2009. Exposure based waiving: the application of the toxicological threshold of concern (TTC) to inhalation exposure for aerosol ingredients in consumer products. *Food Chem. Toxicol.* 47 (6), 1287–1295.
- Cassano, A., Manganaro, A., Martin, T., Young, D., Piclin, N., Pintore, M., Bigoni, D., Benfenati, E., 2010. CAESAR models for developmental toxicity. *Chem. Cent. J.* 4 (Suppl. 1), S4.
- Comiskey, D., Api, A.M., Barratt, C., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Safford, B., Smith, B., Tozer, S., 2015. Novel database for exposure to fragrance ingredients in cosmetics and personal care products. *Regul. Toxicol. Pharmacol.* 72 (3), 660–672.
- Comiskey, D., Api, A.M., Barrett, C., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Rose, J., Safford, B., Smith, B., Tozer, S., 2017. Integrating habits and practices data for soaps, cosmetics and air care products into an existing aggregate exposure model. *Regul. Toxicol. Pharmacol.* 88, 144–156.
- David, R.M., Tyler, T.R., Ouellette, R., Faber, W.D., Banton, M.I., Garman, R.H., Gill, M.W., O'Donoghue, J.L., 1998. Evaluation of subchronic neurotoxicity of n-butyl acetate vapor. *Neurotoxicology* 19 (6), 809–822.
- David, R.M., Tyler, T.R., Ouellette, R., Faber, W.D., Banton, M.I., 2001. Evaluation of subchronic toxicity of n-butyl acetate vapor. *Food Chem. Toxicol.* 39 (8), 877–886.
- ECHA, 2011. Registration Dossier Isobutyl acetate. Retrieved from. <https://echa.europa.eu/registration-dossier/-/registered-dossier/14421/1>.
- ECHA, 2012. Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT Assessment, November 2012 v1.1. <http://echa.europa.eu/>.
- ECHA, 2016. Read-across Assessment Framework (RAAF). www.echa.europa.eu/documents/10162/13628/raaf_en.pdf.
- Frantik, E., Hornychova, M., Horvath, M., 1994. Relative acute neurotoxicity of solvents: isoeffective air concentrations of 48 compounds evaluated in rats and mice. *Environ. Res.* 66 (2), 173–185.
- Haglund, U., Lundberg, I., Zech, L., 1980. Chromosome aberrations and sister chromatid exchanges in Swedish paint industry workers. *Scand. J. Work. Environ. Health* 6 (4), 291–298.
- Henry, B., Foti, C., Alsante, K., 2009. Can light absorption and photostability data be used to assess the photosafety risks in patients for a new drug molecule? *J. Photochem. Photobiol. B Biol.* 96 (1), 57–62.
- IFRA (International Fragrance Association), 2015. Volume of Use Survey, February 2015.
- Iregren, A., Lof, A., Toomingas, A., Wang, Z., 1993. Irritation effects from experimental exposure to n-butyl acetate. *Am. J. Ind. Med.* 24 (6), 727–742.
- Kaneko, T., Wang, P.-Y., Sato, A., 1994. Partition coefficients of some acetate esters and alcohols in water, blood, olive oil, and rat tissues. *Occup. Environ. Med.* 51 (1), 68–72.
- Klecak, G., 1985. The Freund's complete adjuvant test and the open epicutaneous test. In: *Current Problems in Dermatology*, vol. 14. pp. 152–171.
- Eastman Kodak Company, 1996. Submission to EPA, Unpublished.
- Kroes, R., Renwick, A.G., Feron, V., Galli, C.L., Gibney, M., Greim, H., Guy, R.H., Lhuguenot, J.C., van de Sandt, J.J.M., 2007. Application of the threshold of toxicological concern (TTC) to the safety evaluation of cosmetic ingredients. *Food Chem. Toxicol.* 45 (12), 2533–2562.
- Laufersweiler, M.C., Gadagbui, B., Baskerville-Abraham, I.M., Maier, A., Willis, A., et al., 2012. Correlation of chemical structure with reproductive and developmental toxicity as it relates to the use of the threshold of toxicological concern. *Regul. Toxicol. Pharmacol.* 62 (1), 160–182.
- Martinez-Alfaro, M., Carabez-Trejo, A., Gallegos-Corona, M.-A., Pedaza-Aboytes, G., Hernandez-Chan, N.G., Leo-Amador, G.E., 2009. Thinner inhalation effects on oxidative stress and DNA repair in a rat model of abuse. *J. Appl. Toxicol.* 30 (3), 226–232.
- McOmie, W.A., Anderson, H.H., 1949. Comparative Toxicologic Effects of Some Isobutyl Carbinols and Ketones, vol. 2. University California Publications Pharmacology, pp. 217–230 17.
- National Institute for Occupational Safety and Health, 1982. Teratogenic Study of Ethylene and Propylene Oxide and N-Butyl Acetate. Unpublished.
- Nelson, K.W., Ege, J.F., Ross, M., Woodman, L.E., Silverman, L., 1943. Sensory response to certain industrial solvent vapors. *The Journal of Industrial Hygiene and Toxicology. The Journal of Industrial Hygiene and Toxicology* 25 (7), 282–285.
- Norris, J.C., Nachreiner, D.J., Tyler, T.R., Klimisch, H.J., Zimmerman, D.D., 1997. Acute inhalation toxicity studies of n-butyl acetate. *Inhal. Toxicol.* 9 (7), 623–645.
- OECD, 2012. The OECD QSAR Toolbox, v 3.4. <http://www.qsartoolbox.org/>.
- OECD, 2015. Guidance document on the reporting of integrated Approaches to testing and assessment (IATA). ENV/JM/HA(2015)7. <http://www.oecd.org/>.
- Osina, T.M., 1959. Comparative toxicity of propyl propionate and butyl acetate. *Nauch. Trudy Gos. Usovshenst. Vrachei im. S.M. Kirova* 19, 210–218.
- Prah, J.D., Case, M.W., Goldstein, G.M., 1998. 1998 Equivalence of sensory responses to single and mixed volatile organic compounds at equimolar concentrations. *Environ. Health Perspect.* 106 (11), 739–744.
- Querci, V., Mascia, D., 1970. Enzymological and histological findings on liver damage in experimental acetate intoxication. *Med. Lavoro* 61 (10), 524–530.
- Querci, V., Mascia, D., DiPaolo, N., Bassi, G.P., 1970. Acetate pathology. Review of the literature and chemical-experimental studies. *Lav. Um.* 22 (4), 145–167.
- RIFM (Research Institute for Fragrance Materials, Inc), 1973. Report on Human Maximization Studies. Report to RIFM. RIFM Report Number 1802. RIFM, Woodcliff

- Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 1976. Report on Human Maximization Studies. Report to RIFM. RIFM Report Number 1796. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 1987. Report on Human Repeated Insult Patch Test. Report to RIFM. RIFM Report Number 7973. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2013. Report on the Testing of Isobutyl Acetate in the BlueScreen HC Assay (-/+ S9 Metabolic Activation). RIFM Report Number 65245. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2017. Exposure Survey 15 March 2017.
- Rogers, D., Hahn, M., 2010. Extended-connectivity fingerprints. *J. Chem. Inf. Model.* 50 (5), 742–754.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Smith, B., Thomas, R., Tozer, S., 2015. Use of an aggregate exposure model to estimate consumer exposure to fragrance ingredients in personal care and cosmetic products. *Regul. Toxicol. Pharmacol.* 72, 673–682.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Rose, J., Smith, B., Tozer, S., 2017. Application of the expanded Creme RIFM consumer exposure model to fragrance ingredients in cosmetic, personal care and air care products. *Regul. Toxicol. Pharmacol.* 86, 148–156.
- Saillenfait, A.-M., Gallissot, F., Sabate, J.-P., Bourges-Abella, N., Muller, S., 2007. Developmental toxic effects of ethylbenzene or toluene alone and in combination with butyl acetate in rats after inhalation exposure. *J. Appl. Toxicol.* 27 (1), 32–42.
- Salvito, D.T., Senna, R.J., Federle, T.W., 2002. A Framework for prioritizing fragrance materials for aquatic risk assessment. *Environ. Toxicol. Chem.* 21 (6), 1301–1308.
- Sayers, R.R., Schrenk, H.H., Patty, F.A., 1936. Acute response of Guinea pigs to vapors of some new commercial organic compounds. XII. Normal butyl acetate. *Publ. Health Rep.* 51 (36), 1229–1236.
- Schultz, T.W., Amcoff, P., Berggren, E., Gautier, F., Klaric, M., Knight, D.J., Mahony, C., Schwarz, M., White, A., Cronin, M.T.D., 2015. A strategy for structuring and reporting a read-across prediction of toxicity. *Regul. Toxicol. Pharmacol.* 72 (3), 586–601.
- Shen, J., Kromidas, L., Schultz, T., Bhatia, S., 2014. An *in silico* skin absorption model for fragrance materials. *Food Chem. Toxicol.* 74 (12), 164–176.
- Silver, W.L., 1992. Neural and pharmacological basis for nasal irritation. In: *Annals of the New York Academy of Sciences*, vol. 641. pp. 152–163.
- Smyth, H.F., Smyth Jr., H.F., 1928. Inhalation experiments with certain lacquer solvents. *Journal ind. Hyg.* 10 (8), 261–271.
- Smyth Jr., H.F., Carpenter, C.P., Weil, C.S., Pozzani, U.C., 1954. Range-finding toxicity data. List V. *Archives of Ind. Hyg. Archives of Ind. Hyg.* 10, 61–68.
- Smyth Jr., H.F., Carpenter, C.P., Weil, C.S., Pozzani, U.C., Striegel, J.A., 1962. Range-finding toxicity data: list VI. *Am. Ind. Hyg. Assoc. J.* 23, 95–107.
- Union Carbide Co, 1993. Submission to EPA. Unpublished.
- US EPA, 2012a. Estimation Programs Interface Suite for Microsoft Windows, v4.0–v4.11. United States Environmental Protection Agency, Washington, DC, USA.
- US EPA, 2012b. The ECOSAR (ECOLOGICAL Structure Activity Relationship) Class Program for Microsoft Windows, v1.11. United States Environmental Protection Agency, Washington, DC, USA.