




ELSEVIER

Contents lists available at ScienceDirect

Food and Chemical Toxicology

journal homepage: www.elsevier.com/locate/foodchemtox

Short Review

RIFM fragrance ingredient safety assessment, decyl acetate, CAS Registry Number 112-17-4



A.M. Api^a, D. Belsito^b, D. Botelho^a, M. Bruze^c, G.A. Burton Jr.^d, J. Buschmann^e, M.L. Dagli^f, M. Date^a, W. Dekant^g, C. Deodhar^a, M. Francis^a, A.D. Fryer^h, L. Jones^a, K. Joshi^a, S. La Cava^a, A. Lapczynski^a, D.C. Lieblerⁱ, D. O'Brien^a, A. Patel^a, T.M. Penning^j, G. Ritacco^a, J. Romine^a, N. Sadekar^a, D. Salvito^a, T.W. Schultz^k, I.G. Sipes^l, G. Sullivan^{a,*}, Y. Thakkar^a, Y. Tokura^m, S. Tsang^a

^a Research Institute for Fragrance Materials, Inc., 50 Tice Boulevard, Woodcliff Lake, NJ, 07677, USA

^b Member RIFM Expert Panel, Columbia University Medical Center, Department of Dermatology, 161 Fort Washington Ave., New York, NY, 10032, USA

^c Member RIFM Expert Panel, Malmo University Hospital, Department of Occupational & Environmental Dermatology, Sodra Forstadsgatan 101, Entrance 47, Malmo, SE-20502, Sweden

^d Member RIFM Expert Panel, School of Natural Resources & Environment, University of Michigan, Dana Building G110, 440 Church St., Ann Arbor, MI, 48109, USA

^e Member RIFM Expert Panel, Fraunhofer Institute for Toxicology and Experimental Medicine, Nikolai-Fuchs-Strasse 1, 30625, Hannover, Germany

^f Member RIFM Expert Panel, University of Sao Paulo, School of Veterinary Medicine and Animal Science, Department of Pathology, Av. Prof. dr. Orlando Marques de Paiva, 87, Sao Paulo, CEP 05508-900, Brazil

^g Member RIFM Expert Panel, University of Wuerzburg, Department of Toxicology, Versbacher Str. 9, 97078, Wuerzburg, Germany

^h Member RIFM Expert Panel, Oregon Health Science University, 3181 SW Sam Jackson Park Rd., Portland, OR, 97239, USA

ⁱ Member RIFM Expert Panel, Vanderbilt University School of Medicine, Department of Biochemistry, Center in Molecular Toxicology, 638 Robinson Research Building, 2200 Pierce Avenue, Nashville, TN, 37232-0146, USA

^j Member of RIFM Expert Panel, University of Pennsylvania, Perelman School of Medicine, Center of Excellence in Environmental Toxicology, 1316 Biomedical Research Building (BRB) II/III, 421 Curie Boulevard, Philadelphia, PA, 19104-3083, USA

^k Member RIFM Expert Panel, The University of Tennessee, College of Veterinary Medicine, Department of Comparative Medicine, 2407 River Dr., Knoxville, TN, 37996-4500, USA

^l Member RIFM Expert Panel, Department of Pharmacology, University of Arizona, College of Medicine, 1501 North Campbell Avenue, P.O. Box 245050, Tucson, AZ, 85724-5050, USA

^m Member RIFM Expert Panel, The Journal of Dermatological Science (JDS), Editor-in-Chief, Professor and Chairman, Department of Dermatology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, 431-3192, Japan

Version: 032018. This version replaces any previous versions.

Name: Decyl acetate

CAS Registry Number: 112-17-4

Abbreviation/Definition List:

2-Box Model - A RIFM, Inc. proprietary *in silico* tool used to calculate fragrance air exposure concentration

AF - Assessment Factor

BCF - Bioconcentration Factor

Crema RIFM Model - The Crema RIFM Model uses probabilistic (Monte Carlo) simulations to allow full distributions of data sets, providing a more realistic estimate of aggregate exposure to individuals across a population (Comiskey et al., 2015, 2017; Safford et al., 2015, 2017) compared to a deterministic aggregate approach

DEREK - Derek Nexus is an *in silico* tool used to identify structural alerts

DST - Dermal Sensitization Threshold


ECHA - European Chemicals Agency

EU - Europe/European Union

GLP - Good Laboratory Practice

IFRA - The International Fragrance Association

LOEL - Lowest Observable Effect Level



* Corresponding author.

E-mail address: gsullivan@rifm.org (G. Sullivan).

<https://doi.org/10.1016/j.fct.2019.04.026>

Received 17 April 2018; Received in revised form 30 January 2019; Accepted 15 April 2019

Available online 26 April 2019

0278-6915/ © 2019 Elsevier Ltd. All rights reserved.

MOE - Margin of Exposure
MPPD - Multiple-Path Particle Dosimetry. An *in silico* model for inhaled vapors used to simulate fragrance lung deposition
NA - North America
NESIL - No Expected Sensitization Induction Level
NOAEC - No Observed Adverse Effect Concentration
NOAEL - No Observed Adverse Effect Level
NOEC - No Observed Effect Concentration
NOEL - No Observed Effect Level
OECD - Organisation for Economic Co-operation and Development
OECD TG - Organisation for Economic Co-operation and Development Testing Guidelines
PBT - Persistent, Bioaccumulative, and Toxic
PEC/PNEC - Predicted Environmental Concentration/Predicted No Effect Concentration
QRA - Quantitative Risk Assessment
REACH - Registration, Evaluation, Authorisation, and Restriction of Chemicals
RfD - Reference Dose
RIFM - Research Institute for Fragrance Materials
RQ - Risk Quotient
Statistically Significant - Statistically significant difference in reported results as compared to controls with a $p < 0.05$ using appropriate statistical test
TTC - Threshold of Toxicological Concern
UV/Vis spectra - Ultraviolet/Visible spectra
VCF - Volatile Compounds in Food
VoU - Volume of Use **vPvB** - (very) Persistent, (very) Bioaccumulative
WoE - Weight of Evidence

The Expert Panel for Fragrance Safety* concludes that this material is safe under the limits described in this safety assessment.

This safety assessment is based on the RIFM Criteria Document (Api et al., 2015), which should be referred to for clarifications.

Each endpoint discussed in this safety assessment includes the relevant data that were available at the time of writing (version number in the top box is indicative of the date of approval based on a 2-digit month/day/year), both in the RIFM database (consisting of publicly available and proprietary data) and through publicly available information sources (e.g., SciFinder and PubMed). Studies selected for this safety assessment were based on appropriate test criteria, such as acceptable guidelines, sample size, study duration, route of exposure, relevant animal species, most relevant testing endpoints, etc. A key study for each endpoint was selected based on the most conservative endpoint value (e.g., PNEC, NOAEL, LOEL, and NESIL).

*The Expert Panel for Fragrance Safety is an independent body that selects its own members and establishes its own operating procedures. The Expert Panel is comprised of internationally known scientists that provide RIFM with guidance relevant to human health and environmental protection.

Summary: The use of this material under current conditions is supported by existing information.

Decyl acetate was evaluated for genotoxicity, repeated dose toxicity, reproductive toxicity, local respiratory toxicity, phototoxicity/photoallergenicity, skin sensitization, and environmental safety. Data from decyl acetate and read-across analog hexyl propionate (CAS # 2445-76-3) show that decyl acetate is not expected to be genotoxic. Data from decyl acetate and read-across analog 2-butoxyethyl acetate (CAS # 112-07-2) show that this material does not have skin sensitization potential. The repeated dose and developmental endpoints were completed using data from read-across octyl acetate (CAS # 112-14-1), which provided an MOE > 100. The fertility and local respiratory toxicity endpoints were evaluated using the TTC for a Cramer Class I material, and the exposure to decyl acetate is below the TTC (0.03 mg/kg/day and 1.4 mg/day, respectively). The phototoxicity/photoallergenicity endpoint was completed based on UV spectra. The environmental endpoints were evaluated; decyl acetate was found not to be PBT as per the IFRA Environmental Standards, and its risk quotients, based on its current volume of use in Europe and North America (i.e., PEC/PNEC), are < 1.

Human Health Safety Assessment

Genotoxicity: Not genotoxic.

Repeated Dose Toxicity: NOAEL = 500 mg/kg/day.

(RIFM, 2000a; RIFM, 2016)

(Daughtrey et al., 1989b; ECHA Dossier on Octyl acetate)

Reproductive Toxicity: Developmental: NOAEL = 500 mg/kg/day Fertility: No NOAEL available. Exposure is below the TTC.

(Daughtrey et al., 1989a)

Skin Sensitization: Not a concern for skin sensitization.

(Kern et al., 2010)

Phototoxicity/Photoallergenicity: Not phototoxic/photoallergenic.

(UV Spectra, RIFM DB)

Local Respiratory Toxicity: No NOAEC available. Exposure is below the TTC.

Environmental Safety Assessment

Hazard Assessment:

Persistence: Critical Measured Value: 73% (OECD 301F)

RIFM (2012)

Bioaccumulation: Screening-level: 28.5 L/kg

(EPI Suite v4.11; US EPA, 2012a)

Ecotoxicity: Screening-level: 96-h algae EC50: 0.264 mg/L

(ECOSAR; US EPA, 2012b)

Conclusion: Not PBT or vPvB as per IFRA Environmental Standards

Risk Assessment:

Screening-level: PEC/PNEC (North America and Europe) > 1

(RIFM Framework; Salvito et al., 2002)

Critical Ecotoxicity Endpoint: 96-h algae EC50: 0.264 mg/L

(ECOSAR; US EPA, 2012b)

RIFM PNEC is: 0.0264 µg/L

- Revised PEC/PNECs (2015 IFRA VoU): North America and Europe: < 1

1. Identification

- 1. Chemical Name:** Decyl acetate
- 2. CAS Registry Number:** 112-17-4
- 3. Synonyms:** Acetate C-10; Acetic acid, decyl ester; Decanyl acetate; 酢酸デシル(C = 7 ~ 20)エステル; Acetate C10 decylic; Decyl acetate
- 4. Molecular Formula:** C₁₂H₂₄O₂
- 5. Molecular Weight:** 200.32

6. RIFM Number: 483

7. Stereochemistry: Isomer not specified. No stereocenters and no stereoisomers possible.

2. Physical data

- 1. Boiling Point:** 244 °C (FMA), 247.73 °C (US EPA, 2012a)
- 2. Flash Point:** > 200 °F; CC (FMA), > 93 °C (GHS)

3. **Log Kow:** log Pow = 5.6 (RIFM, 2013), 4.79 (US EPA, 2012a)
4. **Melting Point:** 12.58 °C (US EPA, 2012a)
5. **Water Solubility:** 3.517 mg/L (US EPA, 2012a)
6. **Specific Gravity:** 0.864 (FMA)
7. **Vapor Pressure:** 0.0239 mm Hg @ 20 °C (US EPA, 2012a), 0.01 mm Hg @ 20 °C (FMA), 0.0375 mm Hg @ 25 °C (US EPA, 2012a)
8. **UV Spectra:** No significant absorbance between 290 and 700 nm; molar absorption coefficient is below the benchmark (1000 L mol⁻¹ · cm⁻¹)
9. **Appearance/Organoleptic:** A colorless to pale yellow clear liquid with a fresh and fruity odor.*

* <http://www.thegoodscentscompany.com/data/rw1486221.html>, 08/23/17.

3. Exposure

1. **Volume of Use (worldwide band):** 1–10 metric ton per year (IFRA, 2015)
2. **95th Percentile Concentration in Hydroalcohols:** 0.0037% (RIFM, 2016a)
3. **Inhalation Exposure*:** 0.000010 mg/kg/day or 0.00076 mg/day (RIFM, 2016a)
4. **Total Systemic Exposure**:** 0.00013 mg/kg/day (RIFM, 2016a)

*95th percentile calculated exposure derived from concentration survey data in the Creme RIFM aggregate exposure model (Comiskey et al., 2015; Safford et al., 2015; Safford et al., 2017; and Comiskey et al., 2017).

**95th percentile calculated exposure; assumes 100% absorption unless modified by dermal absorption data as reported in Section 4. It is derived from concentration survey data in the Creme RIFM aggregate exposure model and includes exposure via dermal, oral and inhalation routes whenever the fragrance ingredient is used in products that include these routes of exposure (Comiskey et al., 2015; Safford et al., 2015; Safford et al., 2017; and Comiskey et al., 2017).

4. Derivation of systemic absorption

1. **Dermal:** Assumed 100%
2. **Oral:** Assumed 100%
3. **Inhalation:** Assumed 100%

5. Computational toxicology evaluation

1. **Cramer Classification:** Class I, Low

Expert Judgment	Toxtree v 2.6	OECD QSAR Toolbox v 3.2
I	I	I

2. **Analogs Selected:**

- a. **Genotoxicity:** Hexyl propionate (CAS # 2445-76-3)
- b. **Repeated Dose Toxicity:** Octyl acetate (CAS # 112-14-1)
- c. **Reproductive Toxicity:** Octyl acetate (CAS # 112-14-1)
- d. **Skin Sensitization:** 2-Butoxyethyl acetate (CAS # 112-07-2)
- e. **Phototoxicity/Photoallergenicity:** None
- f. **Local Respiratory Toxicity:** None
- g. **Environmental Toxicity:** None

3. **Read-across Justification:** See Appendix below

6. Metabolism

No relevant data available for inclusion in this safety assessment.

7. Natural occurrence (discrete chemical) or composition (NCS)

Decyl acetate is reported to occur in the following foods*:

Apple brandy (<i>Calvados</i>)	Citrus fruits
Apple fresh (<i>Malus</i> species)	Grape brandy
Babaco fruit (<i>Carica pentagona heilborn</i>)	Melon
Blue cheeses	Passion fruit (<i>Passiflora</i> species)
Cardamom (<i>Ellettaria cardamomum maton.</i>)	Plum (<i>Prunus</i> species)
Celery (<i>Apium graveolens</i> L.)	Strawberry (<i>Fragaria</i> species)

*VCF Volatile Compounds in Food: database/Nijssen, L.M.; Ingen-Visscher, C.A. van; Donders, J.J.H. (eds). – Version 15.1 – Zeist (The Netherlands): TNO Triskelion, 1963–2014. A continually updated database containing information on published volatile compounds that have been found in natural (processed) food products. Includes FEMA GRAS and EU-Flavis data.

8. IFRA standard

None.

9. REACH dossier

Dossier available; accessed on 03/12/2018.

10. Summary

10.1. Human health endpoint summaries

10.1.1. Genotoxicity

Based on the current existing data, decyl acetate does not present a concern for genetic toxicity.

10.1.1.1. Risk assessment. The mutagenic activity of decyl acetate has been evaluated in a bacterial reverse mutation assay conducted in compliance with GLP regulations and in accordance with OECD TG 471 using the standard plate incorporation method. *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and TA102 were treated with decyl acetate in ethanol at concentrations up to 5000 µg/plate. No increases in the mean number of revertant colonies were observed at any tested dose in the presence or absence of S9 (RIFM, 2000a). Under the conditions of the study, decyl acetate was not mutagenic in the Ames test.

There are no studies assessing the clastogenic activity of decyl acetate; however, read-across can be made to hexyl propionate (CAS # 2445-76-3; see Section V). The clastogenic activity of hexyl propionate was evaluated in an *in vitro* micronucleus test conducted in compliance with GLP regulations and in accordance with OECD TG 487. Human peripheral blood lymphocytes were treated with hexyl propionate in dimethyl sulfoxide (DMSO) at concentrations up to 1580 µg/mL in the presence and absence of metabolic activation (S9) for 4 and 24 h. Hexyl propionate did not induce binucleated cells with micronuclei when tested up to cytotoxic levels in either non-activated or S9-activated test systems (RIFM, 2016). Under the conditions of the study, hexyl propionate was considered to be non-clastogenic in the *in vitro* micronucleus test and this can be extended to decyl acetate.

Based on the available data decyl acetate does not present a concern for genotoxic potential.

Additional References: RIFM, 1980b.

Literature Search and Risk Assessment Completed on: 07/31/2017.

10.1.2. Repeated dose toxicity

The margin of exposure for decyl acetate is adequate for the repeated dose toxicity endpoint at the current level of use.

10.1.2.1. Risk assessment. There are insufficient repeated dose toxicity data on decyl acetate. Read-across material, octyl acetate (CAS # 112-14-1; see section 5) has sufficient repeated dose toxicity data. Groups of 20 SD rats/sex/dose were gavaged with octyl acetate 5 days per week for 13 weeks at doses of 0 (distilled water), 100, 500, or 1000 mg/kg/day. At week 13, relative liver weights among mid- and high-dose animals were statistically significantly increased as compared to controls. The increase in liver weights was considered to be adaptive due to lack of histopathological evidence (necrosis, fibrosis, inflammation, and steatotic vacuolar degeneration) showing liver cell damage and associated clinical chemistry alterations (Hall et al., 2012). Relative kidney weights among high-dose animals were also statistically significantly increased as compared to control. Gross pathological examinations did not reveal any differences among treated and control group animals. At week 13, microscopic evaluation of the kidneys revealed evidence of mild tubular nephropathy only in the high-dose male rats. The specific findings consisted of an increased incidence of dilated renal tubules (cortical-medullary zone) containing granular casts and regenerative hyperplasia in proximal convoluted tubules. These histopathological findings were not observed in high-dose females or in either sex among mid- and low-dose group animals. Microscopic alterations in kidney among high-dose males were consistent with documented changes of α -2u-globulin nephropathy, which is species-specific to male rats in response to treatment with some hydrocarbons. This effect is not considered a hazard to human health (Lehman-McKeeman and Caudill, 1992; and Lehman-McKeeman et al., 1990). There were no reports of confirmatory staining during histopathological examinations. Thus, the NOAEL was considered to be 500 mg/kg/day based on increased kidney weight among high-dose females (Daughtrey et al., 1989a; also available in ECHA Dossier: Octyl acetate). Therefore, the decyl acetate MOE for the repeated dose toxicity endpoint can be calculated by dividing the octyl acetate NOAEL in mg/kg/day by the total systemic exposure to decyl acetate, 500/0.00013 or 3846154.

In addition, the total systemic exposure to decyl acetate (0.13 μ g/kg/day) is below the TTC (30 μ g/kg bw/day; Kroes et al., 2007) for the repeated dose toxicity endpoint of a Cramer Class I material at the current level of use.

Additional References: None.

Literature Search and Risk Assessment Completed on: 08/16/17.

10.1.3. Reproductive toxicity

The margin of exposure for decyl acetate is adequate for the developmental toxicity endpoint at the current level of use. There are insufficient fertility data on decyl acetate or on any read-across materials. The total systemic exposure to decyl acetate is below the TTC for the reproductive toxicity endpoint of a Cramer Class I material at the current level of use.

10.1.3.1. Risk assessment. There are insufficient developmental toxicity data on decyl acetate. Read-across material, octyl acetate (CAS # 112-14-1; see Section 5) has sufficient developmental toxicity data. A

gavage developmental toxicity study was conducted in Sprague Dawley rats. Groups of 22 mated females/sex/group were gavaged on gestation days (GD) 6–15 with octyl acetate at doses of 0, 100, 500, or 1000 mg/kg neat. Mortality was reported among 2 females from the high-dose group that expired on GDs 10 and 12. Maternal animals in the high-dose group had increased incidence of alopecia, rales, red nasal discharge, and anal-genital staining. Additionally, mean body weights were decreased in high-dose treated maternal rats at GDs 9, 12, 16, and 20, when compared to the control group. Four fetuses from the high-dose group had different types of vertebral anomalies in the form of incomplete ossifications, but those were not statistically significantly different as compared to controls. Visceral examination revealed dilated lateral cerebral ventricles in 2 fetuses in the high-dose group. These anatomical variations were within the historical controls and hence not considered to be toxicologically relevant. Various types of skeletal variations of incomplete ossifications were observed in all groups. The total number of fetuses (litters) with malformations in the control, low-dose, mid-dose, and high-dose groups were 1(1), 1(1), 1(1) and 6 (6), respectively. Thus, the NOAEL for maternal toxicity was considered to be 500 mg/kg/day, based on incidences of clinical observations and decrease in body weights among high-dose group females. The authors of the study report determined the developmental toxicity NOAEL to be 1000 mg/kg/day (Daughtrey et al., 1989a). Since there were anomalies observed in fetuses of the highest dose group, a more conservative NOAEL of 500 mg/kg/day was considered for the developmental toxicity endpoint. Therefore, the decyl acetate MOE for the developmental toxicity endpoint can be calculated by dividing the octyl acetate NOAEL in mg/kg/day by the total systemic exposure to decyl acetate, 500/0.00013 or 3846154.

There are no fertility data on decyl acetate or on any read-across materials that can be used to support the fertility endpoint. The total systemic exposure to decyl acetate (0.13 μ g/kg/day) is below the TTC (30 μ g/kg bw/day; Kroes et al., 2007; Laufersweiler et al., 2012) for the fertility endpoint of a Cramer Class I material at the current level of use.

Additional References: None.

Literature Search and Risk Assessment Completed on: 08/16/17.

10.1.4. Skin sensitization

Based on the existing data and the read-across analog 2-butoxyethyl acetate (CAS # 112-07-2), decyl acetate does not present a concern for skin sensitization.

10.1.4.1. Risk assessment. Limited skin sensitization studies are available for hexyl propionate. Based on the read-across analog 2-butoxyethyl acetate (CAS # 112-07-2; see Section 5), decyl acetate does not present a concern for skin sensitization. The chemical structures of these materials indicate that they would not be expected to react with skin proteins (Toxtree 2.6.13; OECD toolbox v3.4). Read-across material 2-butoxyethyl acetate was found to be negative in the *in vitro* KeratinoSens, U937-CD86, and h-CLAT tests, but positive in Direct Peptide Reactivity Assay (DPRA) (Natsch et al., 2013; Otsubo et al., 2017). However, in a murine local lymph node assay (LLNA), read-across material 2-butoxyethyl acetate was found to be negative up to the maximum tested concentration of 50% which resulted in SI of 1.2 (Kern et al., 2010; Kern et al., 2010). In guinea pigs, a Buehler test did not present reactions indicative of sensitization for the read-across material 2-butoxyethyl acetate (ECHA dossier: 2-butoxyethyl acetate accessed 7/25/17). In a human maximization test, no skin sensitization reactions were observed with 8% or 5520 μ g/cm² decyl acetate in petrolatum (RIFM, 1980a). Based on weight of evidence from structural

analysis, animal and human studies, and from the read-across material 2-butoxyethyl acetate, decyl acetate does not present a concern for skin sensitization.

Additional References: None

Literature Search and Risk Assessment Completed on: 07/25/17.

10.1.5. Phototoxicity/photoallergenicity

Based on the available UV/Vis spectra, decyl acetate would not be expected to present a concern for phototoxicity or photoallergenicity.

10.1.5.1. Risk assessment. There are no phototoxicity studies available for decyl acetate in experimental models. UV/Vis absorption spectra indicate no significant absorption between 290 and 700 nm. The corresponding molar absorption coefficient is well below the benchmark of concern for phototoxicity and photoallergenicity (Henry et al., 2009). Based on lack of absorbance, decyl acetate does not present a concern for phototoxicity or photoallergenicity.

10.1.5.2. UV spectra analysis. UV/Vis absorption spectra (OECD TG 101) were obtained. The spectra indicate no significant absorbance in the range of 290–700 nm. The molar absorption coefficient is below the benchmark of concern for phototoxic effects, $1000 \text{ L mol}^{-1} \cdot \text{cm}^{-1}$ (Henry et al., 2009).

Additional References: None.

Literature Search and Risk Assessment Completed On: 7/12/17.

10.1.6. Local Respiratory Toxicity

The margin of exposure could not be calculated due to lack of appropriate data. The exposure level for decyl acetate is below the Cramer Class I TTC value for inhalation exposure local effects.

10.1.6.1. Risk assessment. There are no inhalation data available on decyl acetate. Based on the Creme RIFM model, the inhalation exposure is 0.00076 mg/day. This exposure is 1842 times lower than the Cramer Class I TTC value of 1.4 mg/day (based on human lung weight of 650 g; Carthew et al., 2009); therefore, the exposure at the current level of use is deemed safe.

Additional References:

Smyth et al., 1969.

Literature Search and Risk Assessment Completed On: 01/09/18.

10.2. Environmental endpoint summary

10.2.1. Screening-level assessment

A screening level risk assessment of decyl acetate was performed following the RIFM Environmental Framework (Salvito et al., 2002), which provides 3 tiered levels of screening for aquatic risk. In Tier 1, only the material's regional VoU, its log K_{OW} , and its molecular weight are needed to estimate a conservative risk quotient (RQ), expressed as the ratio Predicted Environmental Concentration/Predicted No Effect Concentration (PEC/PNEC). A general QSAR with a high uncertainty factor applied is used to predict fish toxicity, as discussed in Salvito et al. (2002). In Tier 2, the RQ is refined by applying a lower uncertainty factor to the PNEC using the ECOSAR model (US EPA, 2012b), which provides chemical class-specific ecotoxicity estimates. Finally, if necessary, Tier 3 is conducted using measured biodegradation and ecotoxicity data to refine the RQ, thus allowing for lower PNEC

uncertainty factors. The data for calculating the PEC and PNEC for this safety assessment are provided in the table below. For the PEC, the range from the most recent IFRA Volume of Use Survey is reviewed. The PEC is then calculated using the actual regional tonnage, not the extremes of the range. Following the RIFM Environmental Framework, decyl acetate was identified as a fragrance material with the potential to present a possible risk to the aquatic environment (i.e., its screening level PEC/PNEC > 1).

A screening-level hazard assessment using EPI Suite v4.11 (US EPA, 2012a) did not identify decyl acetate as possibly being either persistent or bioaccumulative based on its structure and physical-chemical properties. This screening-level hazard assessment considers the potential for a material to be persistent *and* bioaccumulative *and* toxic, or very persistent *and* very bioaccumulative as defined in the Criteria Document (Api et al., 2015). As noted in the Criteria Document, the screening criteria applied are the same as those used in the EU for REACH (ECHA, 2012). For persistence, if the EPI Suite model BIOWIN 3 predicts a value < 2.2 and either BIOWIN 2 or BIOWIN 6 predicts a value < 0.5, then the material is considered potentially persistent. A material would be considered potentially bioaccumulative if the EPI Suite model BCFBAF predicts a fish BCF $\geq 2000 \text{ L/kg}$. Ecotoxicity is determined in the above screening-level risk assessment. If, based on these model outputs (Step 1), additional assessment is required, a WoE-based review is then performed (Step 2). This review considers available data on the material's physical-chemical properties, environmental fate (e.g., OECD Guideline biodegradation studies or die-away studies), fish bioaccumulation, and higher-tier model outputs (e.g., US EPA's BIOWIN and BCFBAF found in EPI Suite v4.11). Data on persistence and bioaccumulation are reported below and summarized in the Environmental Safety Assessment section prior to Section 1.

10.2.2. Risk assessment

Based on current Volume of Use (2015), decyl acetate presents a risk to the aquatic compartment in the screening level assessment.

10.2.2.1. Biodegradation. RIFM, 2000b: Ready biodegradability of the test material was determined in a Closed Bottle test according to the OECD 301D method. Maximum biodegradation after 28 days was 44%. RIFM, 2012: The ready biodegradability of the test material was evaluated in a Manometric Respirometry Test according to the OECD 301F method. Under the conditions of the study, biodegradation of 73% was observed.

10.2.2.2. Ecotoxicity. RIFM, 2000c: A *Daphnia magna* immobilization study was conducted according to the OECD 202 method under semi-static conditions with filled and sealed test vessels. The 48-h EC50 based on time weighted mean measured concentration was reported to be 1.2 mg/L.

10.2.2.3. Other available data. Decyl acetate has been pre-registered for REACH with no additional data at this time.

10.2.3. Risk assessment refinement

Since decyl acetate has passed the screening criteria, measured data is included for completeness only and has not been used in PNEC derivation.

Ecotoxicological data and PNEC derivation (all endpoints reported in mg/L; PNECs in $\mu\text{g/L}$).

Endpoints used to calculate PNEC are underlined.

	LC50 (Fish) (mg/L)	EC50 (<i>Daphnia</i>) (mg/L)	EC50 (Algae) (mg/L)	AF	PNEC (µg/L)	Chemical Class
RIFM Framework Screening Level (Tier 1)	<u>0.1994</u>			1,000,000	0.0001994	
ECOSAR Acute Endpoints (Tier 2) Ver 1.11	0.626	0.965	<u>0.264</u>	10,000	0.0264	Esters
ECOSAR Acute Endpoints (Tier 2) Ver 1.11	0.510	0.375	0.812			Neutral Organic

Exposure information and PEC calculation (following RIFM Environmental Framework: [Salvito et al., 2002](#)).

Exposure	Europe	North America
Log K_{ow} used	5.6	5.6
Biodegradation Factor Used	1	1
Dilution Factor	3	3
Regional Volume of Use Tonnage Band	1–10	< 1
Risk Characterization: PEC/PNEC	< 1	< 1

Based on available data, the RQ for this material is < 1. No further assessment is necessary.

The RIFM PNEC is 0.0264 µg/. The revised PEC/PNECs for EU and NA are < 1; therefore, the material does not present a risk to the aquatic environment at the current reported volumes of use.

Literature Search and Risk Assessment Completed On: 8/1/17.

11. Literature Search*

- **RIFM Database:** Target, Fragrance Structure Activity Group materials, other references, JECFA, CIR, SIDS
- **ECHA:** <http://echa.europa.eu/>
- **NTP:** <https://ntp.niehs.nih.gov/>
- **OECD Toolbox**

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fct.2019.04.026>.

Appendix

Read-across Justification

Methods

The read-across analogs were identified following the strategy for structuring and reporting a read-across prediction of toxicity described in [Schultz et al. \(2015\)](#). The strategy is also consistent with the guidance provided by OECD within Integrated Approaches for Testing and Assessment ([OECD, 2015](#)) and the European Chemical Agency read-across assessment framework ([ECHA, 2016](#)).

- **SciFinder:** <https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf>
- **PubMed:** <http://www.ncbi.nlm.nih.gov/pubmed>
- **TOXNET:** <http://toxnet.nlm.nih.gov/>
- **IARC:** <http://monographs.iarc.fr>
- **OECD SIDS:** <http://webnet.oecd.org/hpv/ui/Default.aspx>
- **EPA ACToR:** <https://actor.epa.gov/actor/home.xhtml>
- **US EPA HPVIS:** https://ofmpub.epa.gov/opthpv/public_search_publicdetails?submission_id=24959241&ShowComments=Yes&sqlstr=null&recordcount=0&User_title=DetailQuery%20Results&EndPointRpt=Y#submission
- **Japanese NITE:** <http://www.safe.nite.go.jp/english/db.html>
- **Japan Existing Chemical Data Base (JECDB):** http://dra4.nihs.go.jp/mhlw_data/jsp/SearchPageENG.jsp
- **Google:** <https://www.google.com>
- **ChemIDplus:** <https://chem.nlm.nih.gov/chemidplus/>




Search keywords: CAS number and/or material names.

*Information sources outside of RIFM's database are noted as appropriate in the safety assessment. This is not an exhaustive list.

Conflicts of interest

The authors declare that they have no conflicts of interest.

- First, materials were clustered based on their structure similarity. Second, data availability and data quality on the selected cluster was examined. Third, appropriate read-across analogs from the cluster were confirmed by expert judgment.
- Tanimoto structure similarity scores were calculated using FCFC4 fingerprints (Rogers and Hahn, 2010).
- The physical–chemical properties of the target substance and the read-across analogs were calculated using EPI Suite v4.11 (US EPA, 2012a).
- J_{\max} values were calculated using RIFM's skin absorption model (SAM). The parameters were calculated using the consensus model (Shen et al., 2014).
- DNA binding, mutagenicity, genotoxicity alerts, and oncologic classification predictions were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- ER binding and repeat dose categorization were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- Developmental toxicity was predicted using CAESAR v2.1.7 (Cassano et al., 2010), and skin sensitization was predicted using Toxtree 2.6.13.
- Protein binding was predicted using OECD QSAR Toolbox v3.4 (OECD, 2012).
- The major metabolites for the target and read-across analogs were determined and evaluated using OECD QSAR Toolbox v3.4 (OECD, 2012).

	Target Material	Read-across Material	Read-across Material	Read-across Material
Principal Name	Decyl acetate	Hexyl propionate	Octyl acetate	2-Butoxyethyl acetate
CAS No.	112-17-4	2445-76-3	112-14-1	112-07-2
Structure				
Similarity (Tanimoto Score)		0.76	0.93	0.51
Read-across Endpoint		<ul style="list-style-type: none"> • Genotoxicity 	<ul style="list-style-type: none"> • Repeated Dose toxicity • Developmental toxicity 	<ul style="list-style-type: none"> • Skin sensitization
Molecular Formula	C ₁₂ H ₂₄ O ₂	C ₉ H ₁₈ O ₂	C ₁₀ H ₂₀ O ₂	C ₈ H ₁₆ O ₃
Molecular Weight	200.32	158.24	172.27	160.21
Melting Point (°C, EPI Suite)	12.58	−20.94	−9.50	−15.23
Boiling Point (°C, EPI Suite)	247.73	190.83	210.70	191.62
Vapor Pressure (Pa @ 25 °C, EPI Suite)	5	79	29.1	71.5
Log Kow (KOWWIN v1.68 in EPI Suite)	4.79	3.32	3.81	1.57
Water Solubility (mg/L, @ 25 °C, WSKOW v1.42 in EPI Suite)	3.517	101.9	33.39	3103
J_{\max} (mg/cm ² /h, SAM)	7.603	78.149	33.5	26.22
Henry's Law (Pa·m ³ /mol, Bond Method, EPI Suite)	2.28E+002	9.73E+001	1.29E+002	6.46E-001
Genotoxicity				
DNA Binding (OASIS v1.4, QSAR Toolbox v3.4)	<ul style="list-style-type: none"> • Schiff base formation • Nucleophilic attack • Acylation • No alert found 	<ul style="list-style-type: none"> • No alert found 		
DNA Binding (OECD QSAR Toolbox v3.4)		<ul style="list-style-type: none"> • No alert found 		
Carcinogenicity (ISS)	<ul style="list-style-type: none"> • Non-carcinogen (low reliability) • No alert found • No alert found • No alert found • No alert found 	<ul style="list-style-type: none"> • Non-carcinogen (low reliability) • No alert found • No alert found • No alert found • No alert found 		
DNA Binding (Ames, MN, CA, OASIS v1.1)	<ul style="list-style-type: none"> • No alert found 	<ul style="list-style-type: none"> • No alert found 		
<i>In Vitro</i> Mutagenicity (Ames, ISS)	<ul style="list-style-type: none"> • No alert found 	<ul style="list-style-type: none"> • No alert found 		
<i>In Vivo</i> Mutagenicity (Micronucleus, ISS)	<ul style="list-style-type: none"> • No alert found 	<ul style="list-style-type: none"> • No alert found 		
Oncologic Classification	<ul style="list-style-type: none"> • Not classified 	<ul style="list-style-type: none"> • Not classified 		
Repeated Dose Toxicity			<ul style="list-style-type: none"> • Not categorized 	
Repeated Dose (HESS)	<ul style="list-style-type: none"> • Not categorized 			
Developmental Toxicity				
ER Binding (OECD QSAR Toolbox v3.4)	<ul style="list-style-type: none"> • Non binder, non cyclic structure 		<ul style="list-style-type: none"> • Non binder, non cyclic structure 	
Developmental Toxicity (CAESAR v2.1.6)	<ul style="list-style-type: none"> • Non-toxicant (moderate reliability) 		<ul style="list-style-type: none"> • Non-toxicant (low reliability) 	
Skin Sensitization				
Protein Binding (OASIS v1.1)	<ul style="list-style-type: none"> • No alert found 			<ul style="list-style-type: none"> • No alert found
Protein Binding (OECD)	<ul style="list-style-type: none"> • No alert found 			<ul style="list-style-type: none"> • No alert found
Protein Binding Potency	<ul style="list-style-type: none"> • Not possible to classify 			<ul style="list-style-type: none"> • Not possible to classify
Protein Binding Alerts for Skin Sensitization (OASIS v1.1)	<ul style="list-style-type: none"> • No alert found 			<ul style="list-style-type: none"> • No alert found
Skin Sensitization Reactivity Domains (Toxtree v2.6.13)	<ul style="list-style-type: none"> • No alert found 			<ul style="list-style-type: none"> • No alert found
Local Respiratory Toxicity				
Respiratory Sensitization (OECD QSAR Toolbox v3.4)	<ul style="list-style-type: none"> • No alert found 			
Metabolism				
Rat Liver S9 Metabolism Simulator and Structural Alerts for Metabolites (OECD QSAR Toolbox v3.4)	See Supplemental Data 1	See Supplemental Data 2	See Supplemental Data 3	See Supplemental Data 4

Summary

There are insufficient toxicity data on decyl acetate (CAS # 112-17-4). Hence, *in silico* evaluation was conducted to determine read-across analogs for this material. Based on structural similarity, reactivity, metabolism, physical–chemical properties, and expert judgment, hexyl propionate (CAS # 2445-76-3), octyl acetate (CAS # 112-14-1), and 2-butoxyethyl acetate (CAS # 112-07-2) were identified as read-across materials with sufficient data for toxicological evaluation.

Conclusions

- Hexyl propionate (CAS # 2445-76-3) was used as a read-across analog for the target material decyl acetate (CAS # 112-17-4) for the genotoxicity endpoint.
 - o The target substance and the read-across analog are structurally similar and belong to the class of aliphatic esters.
 - o The target substance and the read-across analog share a straight chain primary alcohol fragment.
 - o The key difference between the target substance and the read-across analog is that the target substance has a C10 alcohol fragment attached to the acetyl moiety, whereas the read-across analog has a C6 alcohol fragment attached to the propionate moiety. This structural difference is toxicologically insignificant.
 - o Similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score is mainly driven by the straight chain primary alcohol fragment. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
 - o The physical–chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
 - o Differences are predicted for J_{\max} , which estimates skin absorption. $J_{\max} \leq 40\%$ for the target substance and $\leq 80\%$ for the read-across analog. While the percentage of skin absorption estimated from J_{\max} indicates exposure to the substance, it does not represent hazard or toxicity. This parameter provides context to assess the impact of bioavailability on toxicity comparisons between the materials evaluated.
 - o According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
 - o The target compound is given an alert of Schiff base formation by the DNA binding model within OASIS. Other genotoxicity alerts for the target compound are negative. The read-across analog does not have such an alert. The data described for the read-across analog in the genotoxicity section confirms that the read-across material does not pose a concern for genetic toxicity. Therefore, based on structure similarity between the read-across analog and the target substance and the data described for the read-across analog, this alert for the target substance will be superseded by the availability of data for read-across analog.
 - o The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
 - o The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.
- Octyl acetate (CAS # 112-14-1) was used as a read-across analog for the target material decyl acetate (CAS # 112-17-4) for the repeated dose toxicity and developmental toxicity endpoints.
 - o The target substance and the read-across analog are structurally similar and belong to the class of aliphatic esters.
 - o The target substance and the read-across analog share a straight chain primary alcohol fragment.
 - o The key difference between the target substance and the read-across analog is that the target substance has a C10 alcohol fragment, whereas the read-across analog has a C8 alcohol fragment. This structural difference is toxicologically insignificant.
 - o Similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score is mainly driven by the straight chain primary alcohol fragment. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
 - o The physical–chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
 - o Differences are predicted for J_{\max} , which estimates skin absorption. $J_{\max} \leq 40\%$ for the target substance and $\leq 80\%$ for the read-across analog. While the percentage of skin absorption estimated from J_{\max} indicates exposure to the substance, it does not represent hazard or toxicity. This parameter provides context to assess the impact of bioavailability on toxicity comparisons between the materials evaluated.
 - o According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
 - o The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
 - o The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.
- 2-Butoxyethyl acetate (CAS # 112-07-2) was used as a read-across analog for the target material decyl acetate (CAS # 112-17-4) for the skin sensitization endpoint.
 - o The target substance and the read-across analog are structurally similar and belong to the class of aliphatic esters.
 - o The target substance and the read-across analog share a straight chain primary alcohol fragment.
 - o The key difference between the target substance and the read-across analog is that the target substance has a C10 alcohol fragment, and the read-across analog has a C7 alcohol fragment. The read-across analog has an additional inert ether linkage in the alcohol fragment. This structural difference is toxicologically insignificant.
 - o Similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score is mainly driven by the straight chain primary alcohol fragment. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
 - o The physical–chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
 - o Differences are predicted for J_{\max} , which estimates skin absorption. $J_{\max} \leq 40\%$ for the target substance and $\leq 80\%$ for the read-across analog. While the percentage of skin absorption estimated from J_{\max} indicates exposure to the substance, it does not represent hazard or toxicity. This parameter provides context to assess the impact of bioavailability on toxicity comparisons between the materials evaluated.
 - o According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
 - o The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
 - o The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.

References

- Api, A.M., Belsito, D., Bruze, M., Cadby, P., Calow, P., Dagli, M.L., Dekant, W., Ellis, G., Fryer, A.D., Fukayama, M., Griem, P., Hickey, C., Kromidas, L., Lalko, J.F., Liebler, D.C., Miyachi, Y., Politano, V.T., Renskers, K., Ritacco, G., Salvito, D., Schultz, T.W., Sipes, I.G., Smith, B., Vitale, D., Wilcox, D.K., 2015. Criteria for the research Institute for fragrance materials, Inc. (RIFM) safety evaluation process for fragrance ingredients. *Food Chem. Toxicol.* 82, S1–S19.
- Carthew, P., Clapp, C., Gutsell, S., 2009. Exposure based waiving: the application of the

- toxicological threshold of concern (TTC) to inhalation exposure for aerosol ingredients in consumer products. *Food Chem. Toxicol.* 47 (6), 1287–1295.
- Cassano, A., Manganaro, A., Martin, T., Young, D., Piclin, N., Pintore, M., Bigoni, D., Benfenati, E., 2010. CAESAR models for developmental toxicity. *Chem. Cent. J.* 4 (Suppl. 1), S4.
- Comiskey, D., Api, A.M., Barratt, C., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Safford, B., Smith, B., Tozer, S., 2015. Novel database for exposure to fragrance ingredients in cosmetics and personal care products. *Regul. Toxicol. Pharmacol.* 72 (3), 660–672.
- Comiskey, D., Api, A.M., Barrett, C., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Rose, J., Safford, B., Smith, B., Tozer, S., 2017. Integrating habits and practices data for soaps, cosmetics and air care products into an existing aggregate exposure model. *Regul. Toxicol. Pharmacol.* 88, 144–156.
- Daughtrey, W.C., Eutermoser, M., Thompson, S.W., Biles, R.W., 1989a. A subchronic toxicity study of octyl acetate in rats. *Fundam. Appl. Toxicol.* 12, 313–320.
- Daughtrey, W.C., Wier, P.J., Traul, K.A., Biles, R.W., Egan, G.F., 1989b. Evaluation of the teratogenic potential of octyl acetate in rats. *Fundam. Appl. Toxicol.* 13 (2), 303–309.
- ECHA, 2012. Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT Assessment, November 2012 v1.1. <http://echa.europa.eu/>.
- ECHA, 2016. Read-across Assessment Framework (RAAF). Retrieved from www.echa.europa.eu/documents/10162/13628/raaf_en.pdf.
- Hall, A.P., Elcombe, C.R., Foster, J.R., Harada, T., Kaufmann, W., et al., 2012. Liver hypertrophy: a review of adaptive (adverse and non-adverse) changes—conclusions from the 3rd International ESTP expert workshop. *Toxicol. Pathol.* 40 (7), 971–974.
- Henry, B., Foti, C., Alsante, K., 2009. Can light absorption and photostability data be used to assess the photosafety risks in patients for a new drug molecule? *J. Photochem. Photobiol. B Biol.* 96 (1), 57–62.
- IFRA (International Fragrance Association), 2015. Volume of Use Survey, February 2015.
- Kern, P.S., Gerberick, G.F., Ryan, C.A., Kimber, I., Aptula, A., Basketter, D.A., 2010. Local lymph node data for the evaluation of skin sensitization alternatives: a second compilation. *Dermatitis* 21 (1), 8–32.
- Kroes, R., Renwick, A.G., Feron, V., Galli, C.L., Gibney, M., Greim, H., Guy, R.H., Lhuguenot, J.C., van de Sandt, J.J.M., 2007. Application of the threshold of toxicological concern (TTC) to the safety evaluation of cosmetic ingredients. *Food Chem. Toxicol.* 45 (12), 2533–2562.
- Laufersweiler, M.C., Gadagbui, B., Baskerville-Abraham, I.M., Maier, A., Willis, A., et al., 2012. Correlation of chemical structure with reproductive and developmental toxicity as it relates to the use of the threshold of toxicological concern. *Regul. Toxicol. Pharmacol.* 62 (1), 160–182.
- Lehman-McKeeman, L.D., Rivera-Torres, M.I., Caudill, D., 1990. Lysosomal degradation of alpha2u-globulin and alpha2u-globulin-xenobiotic conjugates. *Toxicol. Appl. Pharmacol.* 103 (3), 539–548.
- Lehman-McKeeman, L.D., Caudill, D., 1992. a-2u-globulin is the only member of the lipocalin protein superfamily that binds to hyaline droplet inducing agents. *Toxicol. Appl. Pharmacol.* 116 (2), 170–176.
- Natsch, A., Ryan, C.A., Foertsch, L., Emter, R., Jaworska, J., Gerberick, F., Kern, P., 2013. A dataset on 145 chemicals tested in alternative assays for skin sensitization undergoing prevalidation. *J. Appl. Toxicol.* 33 (11), 1337–1352.
- OECD, 2012. The OECD QSAR Toolbox, V 3.4. Retrieved from <http://www.qsar toolbox.org/>.
- OECD, 2015. Guidance Document on the Reporting of Integrated Approaches to Testing and Assessment (IATA). ENV/JM/HA(2015)7. Retrieved from <http://www.oecd.org/>.
- Otsubo, Y., Nishijo, T., Miyazawa, M., Saito, K., Mizumachi, H., Sakaguchi, H., 2017. Binary test battery with KeratinoSens™ and h-CLAT as part of a bottom-up approach for skin sensitization hazard prediction. *Regul. Toxicol. Pharmacol.* 88, 118–124.
- RIFM (Research Institute for Fragrance Materials, Inc), 1980a. Report on Human Maximization Studies. Report to RIFM. RIFM Report Number 1790. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 1980b. Ames Metabolic Activation Test to Assess the Potential Mutagenic Effect of Decyl Acetate. Unpublished report from Quest International. RIFM report number 45903. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2000a. Mutagenicity Study of Decyl Acetate in the Salmonella typhimurium/mammalian Microsome Reverse Mutation Assay (Ames-Test). Unpublished report from Symrise. RIFM report number 60696. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2000b. Decyl Acetate: Ready Biodegradability Closed Bottle Test. Unpublished report from Symrise. RIFM report number 60697. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2000c. Decyl Acetate: Acute Immobilisation Test (48 H) to Daphnia Magna STRAUS. Unpublished report from Symrise. RIFM report number 60698. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2012. Ready Biodegradability of Decyl Acetate (Acetate C10 Decylic). Unpublished report from Givaudan. RIFM report number 63205. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2013. Partition Coefficient N-Octanol/water of Decyl Acetate (Acetate C10 Decylic). Unpublished report from Givaudan. RIFM report number 65185. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2016. Hexyl Propionate: in Vitro Mammalian Cell Micronucleus Assay in Human Peripheral Blood Lymphocytes (HPBL). RIFM Report Number 70466. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2016a. Exposure Survey 09, January 2016.
- Rogers, D., Hahn, M., 2010. Extended-connectivity fingerprints. *J. Chem. Inf. Model.* 50 (5), 742–754.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Smith, B., Thomas, R., Tozer, S., 2015. Use of an aggregate exposure model to estimate consumer exposure to fragrance ingredients in personal care and cosmetic products. *Regul. Toxicol. Pharmacol.* 72, 673–682.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Rose, J., Smith, B., Tozer, S., 2017. Application of the expanded Creme RIFM consumer exposure model to fragrance ingredients in cosmetic, personal care and air care products. *Regul. Toxicol. Pharmacol.* 86, 148–156.
- Salvito, D.T., Senna, R.J., Federle, T.W., 2002. A Framework for prioritizing fragrance materials for aquatic risk assessment. *Environ. Toxicol. Chem.* 21 (6), 1301–1308.
- Schultz, T.W., Amcoff, P., Berggren, E., Gautier, F., Klaric, M., Knight, D.J., Mahony, C., Schwarz, M., White, A., Cronin, M.T.D., 2015. A strategy for structuring and reporting a read-across prediction of toxicity. *Regul. Toxicol. Pharmacol.* 72 (3), 586–601.
- Shen, J., Kromidas, L., Schultz, T., Bhatia, S., 2014. An *in silico* skin absorption model for fragrance materials. *Food Chem. Toxicol.* 74 (12), 164–176.
- Smyth Jr., H.F., Carpenter, C.P., Weil, C.S., Pozzani, U.C., Striegel, J.A., Nycum, J.S., 1969. Range-finding toxicity data: list VII. *Am. Ind. Hyg. Assoc. J.* 30 (5), 470–476.
- US EPA, 2012a. Estimation Programs Interface Suite for Microsoft Windows, v4.0–v4.11. United States Environmental Protection Agency, Washington, DC, USA.
- US EPA, 2012b. The ECOSAR (ECOLOGICAL Structure Activity Relationship) Class Program for Microsoft Windows, v1.11. United States Environmental Protection Agency, Washington, DC, USA.