



ELSEVIER

Contents lists available at ScienceDirect

Food and Chemical Toxicology

journal homepage: www.elsevier.com/locate/foodchemtox

Short Review

RIFM fragrance ingredient safety assessment, methyl 3,7-dimethyl-6-octenoate, CAS Registry Number 2270-60-2



A.M. Api^a, D. Belsito^b, D. Botelho^a, M. Bruze^c, G.A. Burton Jr.^d, J. Buschmann^e, M.L. Dagli^f, M. Date^a, W. Dekant^g, C. Deodhar^a, M. Francis^a, A.D. Fryer^h, L. Jones^a, K. Joshi^a, S. La Cava^a, A. Lapczynski^a, D.C. Lieblerⁱ, D. O'Brien^a, A. Patel^a, T.M. Penning^j, G. Ritacco^a, J. Romine^a, N. Sadekar^a, D. Salvito^a, T.W. Schultz^k, I.G. Sipes^l, G. Sullivan^{a,*}, Y. Thakkar^a, Y. Tokura^m, S. Tsang^a

^a Research Institute for Fragrance Materials, Inc., 50 Tice Boulevard, Woodcliff Lake, NJ, 07677, USA

^b Member RIFM Expert Panel, Columbia University Medical Center, Department of Dermatology, 161 Fort Washington Ave., New York, NY, 10032, USA

^c Member RIFM Expert Panel, Malmo University Hospital, Department of Occupational & Environmental Dermatology, Sodra Forstadsgratan 101, Entrance 47, Malmo, SE-20502, Sweden

^d Member RIFM Expert Panel, School of Natural Resources & Environment, University of Michigan, Dana Building G110, 440 Church St., Ann Arbor, MI, 48109, USA

^e Member RIFM Expert Panel, Fraunhofer Institute for Toxicology and Experimental Medicine, Nikolai-Fuchs-Strasse 1, 30625, Hannover, Germany

^f Member RIFM Expert Panel, University of Sao Paulo, School of Veterinary Medicine and Animal Science, Department of Pathology, Av. Prof. dr. Orlando Marques de Paiva, 87, Sao Paulo, CEP 05508-900, Brazil

^g Member RIFM Expert Panel, University of Wuerzburg, Department of Toxicology, Versbacher Str. 9, 97078, Wuerzburg, Germany

^h Member RIFM Expert Panel, Oregon Health Science University, 3181 SW Sam Jackson Park Rd., Portland, OR, 97239, USA

ⁱ Member RIFM Expert Panel, Vanderbilt University School of Medicine, Department of Biochemistry, Center in Molecular Toxicology, 638 Robinson Research Building, 2200 Pierce Avenue, Nashville, TN, 37232-0146, USA

^j Member of RIFM Expert Panel, University of Pennsylvania, Perelman School of Medicine, Center of Excellence in Environmental Toxicology, 1316 Biomedical Research Building (BRB) II/III, 421 Curie Boulevard, Philadelphia, PA, 19104-3083, USA

^k Member RIFM Expert Panel, The University of Tennessee, College of Veterinary Medicine, Department of Comparative Medicine, 2407 River Dr., Knoxville, TN 37996-4500, USA

^l Member RIFM Expert Panel, Department of Pharmacology, University of Arizona, College of Medicine, 1501 North Campbell Avenue, P.O. Box 245050, Tucson, AZ, 85724-5050, USA

^m Member RIFM Expert Panel, The Journal of Dermatological Science (JDS), Editor-in-Chief, Professor and Chairman, Department of Dermatology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, 431-3192, Japan

Version: 080218. This version replaces any previous versions.

Name: Methyl 3,7-dimethyl-6-octenoate

CAS Registry Number: 2270-60-2

Abbreviation/Definition List:

2-Box Model - A RIFM, Inc. proprietary *in silico* tool used to calculate fragrance air exposure concentration

AF - Assessment Factor

BCF - Bioconcentration Factor

Crete RIFM Model - The Crete RIFM Model uses probabilistic (Monte Carlo) simulations to allow full distributions of data sets, providing a more realistic estimate of aggregate exposure to individuals across a population (Comiskey et al., 2015, 2017; Safford et al., 2015a, 2017) compared to a deterministic aggregate approach

DEREK - Derek Nexus is an *in silico* tool used to identify structural alerts

DST - Dermal Sensitization Threshold

ECHA - European Chemicals Agency

EU - Europe/European Union

GLP - Good Laboratory Practice


IFRA - The International Fragrance Association

LOEL - Lowest Observable Effect Level

MOE - Margin of Exposure

MPPD - Multiple-Path Particle Dosimetry. An *in silico* model for inhaled vapors used to simulate fragrance lung deposition

NA - North America



* Corresponding author.

E-mail address: gsullivan@rifm.org (G. Sullivan).

<https://doi.org/10.1016/j.fct.2019.110616>

Received 23 August 2018; Received in revised form 22 May 2019; Accepted 19 June 2019

Available online 22 June 2019

0278-6915/ © 2019 Elsevier Ltd. All rights reserved.

NESIL - No Expected Sensitization Induction Level
NOAEC - No Observed Adverse Effect Concentration
NOAEL - No Observed Adverse Effect Level
NOEC - No Observed Effect Concentration
NOEL - No Observed Effect Level
OECD - Organisation for Economic Co-operation and Development
OECD TG - Organisation for Economic Co-operation and Development Testing Guidelines
PBT - Persistent, Bioaccumulative, and Toxic
PEC/PNEC - Predicted Environmental Concentration/Predicted No Effect Concentration
QRA - Quantitative Risk Assessment
REACH - Registration, Evaluation, Authorisation, and Restriction of Chemicals
RfD - Reference Dose
RIFM - Research Institute for Fragrance Materials
RQ - Risk Quotient
Statistically Significant - statistically significant difference in reported results as compared to controls with a $p < 0.05$ using appropriate statistical test.
TTC - Threshold of Toxicological Concern
UV/Vis Spectra - Ultraviolet/Visible Spectra
VCF - Volatile Compounds in Food
VoU - Volume of Use **vPvB** - (very) Persistent, (very) Bioaccumulative
WoE - Weight of Evidence

The Expert Panel for Fragrance Safety* concludes that this material is safe as described in this safety assessment.

This safety assessment is based on the RIFM Criteria Document (Api et al., 2015), which should be referred to for clarifications.

Each endpoint discussed in this safety assessment includes the relevant data that were available at the time of writing (version number in the top box is indicative of the date of approval based on a 2-digit month/day/year), both in the RIFM database (consisting of publicly available and proprietary data) and through publicly available information sources (e.g., SciFinder and PubMed). Studies selected for this safety assessment were based on appropriate test criteria, such as acceptable guidelines, sample size, study duration, route of exposure, relevant animal species, most relevant testing endpoints, etc. A key study for each endpoint was selected based on the most conservative endpoint value (e.g., PNEC, NOAEL, LOEL, and NESIL).

*The Expert Panel for Fragrance Safety is an independent body that selects its own members and establishes its own operating procedures. The Expert Panel is comprised of internationally known scientists that provide RIFM with guidance relevant to human health and environmental protection.

Summary: The existing information supports the use of this material as described in this safety assessment.

Methyl 3,7-dimethyl-6-octenoate was evaluated for genotoxicity, repeated dose toxicity, reproductive toxicity, local respiratory toxicity, phototoxicity/photoallergenicity, skin sensitization, and environmental safety. Data from read-across analog *trans*-methylgeranate (CAS # 1189-09-9) show that methyl 3,7-dimethyl-6-octenoate is not expected to be genotoxic. The skin sensitization endpoint was completed using the DST for non-reactive materials (900 $\mu\text{g}/\text{cm}^2$); exposure is below the DST. The repeated dose, reproductive, and local respiratory toxicity endpoints were evaluated using the TTC for a Cramer Class I material and the exposure to methyl 3,7-dimethyl-6-octenoate is below the TTC (0.03 mg/kg/day, 0.03 mg/kg/day, and 1.4 mg/day, respectively). The phototoxicity/photoallergenicity endpoints were evaluated based on UV spectra; methyl 3,7-dimethyl-6-octenoate is not expected to be phototoxic/photoallergenic. The environmental endpoints were evaluated; methyl 3,7-dimethyl-6-octenoate was found not to be PBT as per the IFRA Environmental Standards, and its risk quotients, based on its current volume of use in Europe and North America (i.e., PEC/PNEC), are < 1 .

Human Health Safety Assessment

Genotoxicity: Not expected to be genotoxic.

(RIFM, 2014a; RIFM, 2014b)

Repeated Dose Toxicity: No NOAEL available. Exposure is below the TTC.

Reproductive Toxicity: No NOAEL available. Exposure is below the TTC.

Skin Sensitization: No safety concerns at current, declared use levels; exposure is below the DST.

Phototoxicity/Photoallergenicity: Not expected to be phototoxic/photoallergenic.

(UV Spectra, RIFM Database)

Local Respiratory Toxicity: No NOAEC available. Exposure is below the TTC.

Environmental Safety Assessment

Hazard Assessment:

Persistence: Screening-level: 2.9 (BIOWIN 3)

(EPI Suite v4.11; US EPA, 2012a)

Bioaccumulation: Screening-level: 225 L/kg

(EPI Suite v4.11; US EPA, 2012a)

Ecotoxicity: Screening-level: Fish LC50: 3.93 mg/L

(RIFM Framework; Salvito et al., 2002)

Conclusion: Not PBT or vPvB as per IFRA Environmental Standards

Risk Assessment:

Screening-level: PEC/PNEC (North America and Europe) < 1

(RIFM Framework; Salvito et al., 2002)

Critical Ecotoxicity Endpoint: Fish LC50: 3.93 mg/L

(RIFM Framework; Salvito et al., 2002)

RIFM PNEC is: 0.00393 $\mu\text{g}/\text{L}$

- Revised PEC/PNECs (2015 IFRA VoU): North America and Europe: not applicable; cleared at screening-level

1. Identification

- 1. Chemical Name:** Methyl 3,7-dimethyl-6-octenoate
- 2. CAS Registry Number:** 2270-60-2
- 3. Synonyms:** Methyl citronellate; 6-Octenoic acid, 3,7-dimethyl-, methyl ester; Methyl 3,7-dimethyloct-6-enoate; Methyl 3,7-dimethyl-6-octenoate
- 4. Molecular Formula:** $\text{C}_{11}\text{H}_{20}\text{O}_2$
- 5. Molecular Weight:** 184.28
- 6. RIFM Number:** 6730
- 7. Stereochemistry:** Isomer not specified. One stereocenter and 2 total stereoisomers possible.

2. Physical data

- 1. Boiling Point:** 219.01 $^{\circ}\text{C}$ (EPI Suite)
- 2. Flash Point:** Not Available
- 3. Log K_{ow} :** 4.07 (EPI Suite)
- 4. Melting Point:** 18.46 $^{\circ}\text{C}$ (EPI Suite)
- 5. Water Solubility:** 17.56 mg/L (EPI Suite)
- 6. Specific Gravity:** 0.89700 to 0.90700 @ 25 $^{\circ}\text{C}^*$
- 7. Vapor Pressure:** 0.091 mm Hg @ 20 $^{\circ}\text{C}$ (EPI Suite v4.0), 0.08 mm Hg @ 20 $^{\circ}\text{C}$ (FMA Database), 0.138 mm Hg @ 25 $^{\circ}\text{C}$ (EPI Suite)
- 8. UV Spectra:** No significant absorbance between 290 and 700 nm; molar absorption coefficient is below the benchmark (1000 $\text{L mol}^{-1} \cdot \text{cm}^{-1}$)

9. **Appearance/Organoleptic:** colorless mobile liquid, refreshing, winey-green, fruity and brandy-like, apple-like odor of moderate to poor tenacity (Arctander Volume II 1969)

*<http://www.thegoodscentscompany.com/data/rw1035931.html>, retrieved 12/6/2017.

3. Exposure to fragrance ingredient

1. **Volume of Use (Worldwide Band):** 0.1–1 metric tons per year (IFRA, 2015)
2. **95th Percentile Concentration in Hydroalcoholics:** 0.00026% (RIFM, 2017)
3. **Inhalation Exposure*:** 0.0000007 mg/kg/day or 0.000051 mg/day (RIFM, 2017)
4. **Total Systemic Exposure**:** 0.000018 mg/kg/day (RIFM, 2017)

*95th percentile calculated exposure derived from concentration survey data in the Creme RIFM aggregate exposure model (Comiskey et al., 2015, 2017; Safford et al., 2015a, 2017).

**95th percentile calculated exposure; assumes 100% absorption unless modified by dermal absorption data as reported in Section IV. It is derived from concentration survey data in the Creme RIFM aggregate exposure model and includes exposure via dermal, oral, and inhalation routes whenever the fragrance ingredient is used in products that include these routes of exposure (Comiskey et al., 2015, 2017; Safford et al., 2015a, 2017).

4. Derivation of systemic absorption

1. **Dermal:** Assumed 100%
2. **Oral:** Assumed 100%
3. **Inhalation:** Assumed 100%

5. Computational toxicology evaluation

1. **Cramer Classification:** Class I, Low

Expert Judgment	Toxtree v 2.6	OECD QSAR Toolbox v 3.2
I	I	I

2. Analogs Selected:

- a. **Genotoxicity:** *trans*-methylgeranate (CAS # 1189-09-9)
 - b. **Repeated Dose Toxicity:** None
 - c. **Reproductive Toxicity:** None
 - d. **Skin Sensitization:** None
 - e. **Phototoxicity/Photoallergenicity:** None
 - f. **Local Respiratory Toxicity:** None
 - g. **Environmental Toxicity:** None
3. Read-across Justification: See Appendix below

6. Metabolism

No relevant data available for inclusion in this safety assessment.

7. Natural occurrence (discrete chemical) or composition (NCS)

Methyl 3,7-dimethyl-6-octenoate is reported to occur in the following foods* and in some natural complex substances (NCS):

- Apple Brandy (*Calvados*).
- Lemon Balm (*Melissa officinalis* L.)
- Pepper (*Piper nigrum* L.)

*VCF Volatile Compounds in Food: Database/Nijssen, L.M.; Ingen-Visscher, C.A. van; Donders, J.J.H. (eds). – Version 15.1 – Zeist (The

Netherlands): TNO Triskelion, 1963–2014. A continually updated database containing information on published volatile compounds that have been found in natural (processed) food products. Includes FEMA GRAS and EU-Flavis data.

8. IFRA standard

None.

9. REACH dossier

Pre-registered for 2010, no dossier available as of 08/02/2018.

10. Summary

10.1. Human health endpoint summaries

10.1.1. Genotoxicity

Based on the current existing data, methyl 3,7-dimethyl-6-octenoate does not present a concern for genotoxicity.

10.1.1.1. Risk assessment. Methyl 3,7-dimethyl-6-octenoate was assessed in the BlueScreen assay and found negative for genotoxicity, with and without metabolic activation (RIFM, 2013). BlueScreen is a screening assay that assesses genotoxic stress through human-derived gene expression. Additional assays on a more reactive read-across material were considered to fully assess the potential mutagenic or clastogenic effects on the target material.

There are no data assessing the mutagenic activity of methyl 3,7-dimethyl-6-octenoate; however, read-across can be made to *trans*-methylgeranate (CAS # 1189-09-9; see Section V). The mutagenic activity of *trans*-methylgeranate has been evaluated in a bacterial reverse mutation assay conducted in compliance with GLP regulations and in accordance with OECD TG 471 using the standard plate incorporation method. *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and *Escherichia coli* strain WP2uvrA were treated with *trans*-methylgeranate in dimethyl sulfoxide (DMSO) at concentrations up to 5000 µg/plate. No increases in the mean number of revertant colonies were observed at any tested concentration in the presence or absence of S9 (RIFM, 2014a). Under the conditions of the study, *trans*-methylgeranate was not mutagenic in the Ames test, and this can be extended to methyl 3,7-dimethyl-6-octenoate.

There are no studies assessing the clastogenic activity of methyl 3,7-dimethyl-6-octenoate; however, read-across can be made to *trans*-methylgeranate (CAS # 1189-09-9; see Section V). The clastogenic activity of *trans*-methylgeranate was evaluated in an *in vitro* micronucleus test conducted in compliance with GLP regulations and in accordance with OECD TG 487. Human peripheral blood lymphocytes were treated with *trans*-methylgeranate in DMSO at concentrations up to 1825 µg/mL in the presence and absence of metabolic activation (S9) for 3 h and in the absence of metabolic activation for 24 h. Micronuclei analysis was conducted up to doses producing appropriate cytotoxicity (up to 256 µg/mL) in all the test conditions. *trans*-Methylgeranate did not induce binucleated cells with micronuclei when tested up to cytotoxic levels in either the presence or absence of an S9 activation system (RIFM, 2014b). Under the conditions of the study, *trans*-methylgeranate was considered to be non-clastogenic in the *in vitro* micronucleus test, and this can be extended to methyl 3,7-dimethyl-6-octenoate.

Based on the data available, methyl 3,7-dimethyl-6-octenoate does not present a concern for genotoxic potential.

Additional References: None.

Literature Search and Risk Assessment Completed On: 12/1/2017.

10.1.2. Repeated dose toxicity

There are insufficient repeated dose toxicity data on methyl 3,7-

dimethyl-6-octenoate or any read-across materials. The total systemic exposure to methyl 3,7-dimethyl-6-octenoate is below the TTC for the repeated dose toxicity endpoint of a Cramer Class I material at the current level of use.

10.1.2.1. Risk assessment. There are no repeated dose toxicity data on methyl 3,7-dimethyl-6-octenoate or any read-across materials that can be used to support the repeated dose toxicity endpoint. The total systemic exposure to methyl 3,7-dimethyl-6-octenoate (0.018 µg/kg bw/day) is below the TTC (30 µg/kg bw/day; Kroes et al., 2007) for the repeated dose toxicity endpoint of a Cramer Class I material at the current level of use.

Additional References: None.

Literature Search and Risk Assessment Completed On: 11/22/17.

10.1.3. Reproductive toxicity

There are insufficient reproductive toxicity data on methyl 3,7-dimethyl-6-octenoate or any read-across materials. The total systemic exposure to methyl 3,7-dimethyl-6-octenoate is below the TTC for the reproductive toxicity endpoint of a Cramer Class I material at the current level of use.

10.1.3.1. Risk assessment. There are no reproductive toxicity data on methyl 3,7-dimethyl-6-octenoate or any read-across materials that can be used to support the reproductive toxicity endpoint. The total systemic exposure to methyl 3,7-dimethyl-6-octenoate (0.018 µg/kg bw/day) is below the TTC (30 µg/kg bw/day; Kroes et al., 2007; Laufersweiler et al., 2012) for the reproductive toxicity endpoint of a Cramer Class I material at the current level of use.

Additional References: None.

Literature Search and Risk Assessment Completed On: 11/22/17.

10.1.4. Skin sensitization

Based on the application of DST, methyl 3,7-dimethyl-6-octenoate does not present a safety concern for skin sensitization under the current, declared levels of use.

10.1.4.1. Risk assessment. The chemical structure of this material indicates that it would not be expected to react with skin proteins (Toxtree 2.6.13; OECD toolbox v3.4). No skin sensitization studies are available for methyl 3,7-dimethyl-6-octenoate or read-across materials.

Table 1

Acceptable concentrations for methyl 3,7-dimethyl-6-octenoate that present no appreciable risk for skin sensitization based on non-reactive DST.

IFRA Category ^a	Description of Product Type	Acceptable Concentrations in Finished Products Based on Non-reactive DST	Reported 95th Percentile Concentration in Finished Products
1	Products applied to the lips	0.07%	0.00%
2	Products applied to the axillae	0.02%	0.00% ^b
3	Products applied to the face using fingertips	0.41%	0.00% ^b
4	Fine fragrance products	0.39%	0.00% ^b
5	Products applied to the face and body using the hands (palms), primarily leave-on	0.10%	0.00% ^b
6	Products with oral and lip exposure	0.23%	0.00%
7	Products applied to the hair with some hand contact	0.79%	0.00%
8	Products with significant ano-genital exposure	0.04%	No Data ^c
9	Products with body and hand exposure, primarily rinse-off	0.75%	0.00% ^b
10	Household care products with mostly hand contact	2.70%	0.00% ^b
11	Products with intended skin contact but minimal transfer of fragrance to skin from inert substrate	1.50%	No Data ^c
12	Products not intended for direct skin contact, minimal or insignificant transfer to skin	Not Restricted	0.00% ^b

Note.

^a For a description of the categories, refer to the IFRA/RIFM Information Booklet.

^b Negligible exposure (< 0.01%).

^c Fragrance exposure from these products is very low. These products are not currently in the Creme RIFM Aggregate Exposure Model.

Acting conservatively, due to the insufficient data, the reported exposure was benchmarked utilizing the non-reactive Dermal Sensitization Threshold (DST) of 900 µg/cm² (Safford, 2008; Safford et al., 2011, 2015b; Roberts et al., 2015). The current exposure from the 95th percentile concentration is below the DST for non-reactive materials when evaluated in all QRA categories. Table 1 provides the acceptable concentrations for methyl 3,7-dimethyl-6-octenoate that present no appreciable risk for skin sensitization based on the non-reactive DST. These concentrations are not limits; they represent acceptable concentrations based on the DST approach.

Additional References: None.

Literature Search and Risk Assessment Completed On: 11/16/17.

10.1.5. Phototoxicity/photoallergenicity

Based on the available UV/Vis spectra, methyl 3,7-dimethyl-6-octenoate would not be expected to present a concern for phototoxicity or photoallergenicity.

10.1.5.1. Risk assessment. There are no phototoxicity studies available for methyl 3,7-dimethyl-6-octenoate in experimental models. UV/Vis absorption spectra indicate no significant absorption between 290 and 700 nm. The corresponding molar absorption coefficient is well below the benchmark of concern for phototoxicity and photoallergenicity (Henry et al., 2009). Based on lack of absorbance, methyl 3,7-dimethyl-6-octenoate does not present a concern for phototoxicity or photoallergenicity.

10.1.5.2. UV spectra analysis. UV/Vis absorption spectra (OECD TG 101) were obtained. The spectra indicate no significant absorbance in the range of 290–700 nm. The molar absorption coefficient is below the benchmark of concern for phototoxic effects, 1000 L mol⁻¹ · cm⁻¹ (Henry et al., 2009).

Additional References: None.

Literature Search and Risk Assessment Completed On: 10/20/17.

10.1.6. Local Respiratory Toxicity

The margin of exposure could not be calculated due to lack of appropriate data. The exposure level for methyl 3,7-dimethyl-6-octenoate is below the Cramer Class I TTC value for inhalation exposure local effects.

10.1.6.1. Risk assessment. There are no inhalation data available on methyl 3,7-dimethyl-6-octenoate. Based on the Creme RIFM Model, the inhalation exposure is 0.000051 mg/day. This exposure is 27451 times lower than the Cramer Class I TTC value of 1.4 mg/day (based on human lung weight of 650 g; Carthew et al., 2009); therefore, the exposure at the current level of use is deemed safe.

Additional References: None.

Literature Search and Risk Assessment Completed On: 12/1/2017.

10.2. Environmental endpoint summary

10.2.1. Screening-level assessment

A screening-level risk assessment of methyl 3,7-dimethyl-6-octenoate was performed following the RIFM Environmental Framework (Salvito et al., 2002), which provides 3 tiered levels of screening for aquatic risk. In Tier 1, only the material's regional VoU, its log K_{OW} , and its molecular weight are needed to estimate a conservative risk quotient (RQ), expressed as the ratio Predicted Environmental Concentration/Predicted No Effect Concentration (PEC/PNEC). A general QSAR with a high uncertainty factor applied is used to predict fish toxicity, as discussed in Salvito et al. (2002). In Tier 2, the RQ is refined by applying a lower uncertainty factor to the PNEC using the ECOSAR model (US

available data on the material's physical-chemical properties, environmental fate (e.g., OECD Guideline biodegradation studies or die-away studies), fish bioaccumulation, and higher-tier model outputs (e.g., US EPA's BIOWIN and BCFBAF found in EPI Suite v4.11). Data on persistence and bioaccumulation are reported below and summarized in the Environmental Safety Assessment section prior to Section 1.

10.2.2. Risk assessment

Based on the current Volume of Use (2015), methyl 3,7-dimethyl-6-octenoate does not present a risk to the aquatic compartment in the screening-level assessment.

10.2.2.1. Biodegradation. No data available.

10.2.2.2. Ecotoxicity. No data available.

10.2.2.3. Other available data. Methyl 3,7-dimethyl-6-octenoate has been pre-registered for REACH with no additional data at this time.

10.2.3. Risk assessment refinement

Ecotoxicological data and PNEC derivation (all endpoints reported in mg/L; PNECs in $\mu\text{g/L}$).

Endpoints used to calculate PNEC are underlined.

	LC50 (Fish) (mg/L)	EC50 (<i>Daphnia</i>) (mg/L)	EC50 (Algae) (mg/L)	AF	PNEC ($\mu\text{g/L}$)	Chemical Class
RIFM Framework Screening-level (Tier 1)	<u>3.93</u>			1,000,000	0.00393	

EPA, 2012b), which provides chemical class-specific ecotoxicity estimates. Finally, if necessary, Tier 3 is conducted using measured biodegradation and ecotoxicity data to refine the RQ, thus allowing for lower PNEC uncertainty factors. The data for calculating the PEC and PNEC for this safety assessment are provided in the table below. For the PEC, the range from the most recent IFRA Volume of Use Survey is reviewed. The PEC is then calculated using the actual regional tonnage, not the extremes of the range. Following the RIFM Environmental Framework, methyl 3,7-dimethyl-6-octenoate was identified as a fragrance material with no potential to present a possible risk to the aquatic environment (i.e., its screening-level PEC/PNEC < 1).

A screening-level hazard assessment using EPI Suite v4.11 (US EPA, 2012a) did not identify methyl 3,7-dimethyl-6-octenoate as possibly persistent or bioaccumulative based on its structure and physical-chemical properties. This screening-level hazard assessment considers the potential for a material to be persistent and bioaccumulative and toxic, or very persistent and very bioaccumulative as defined in the Criteria Document (Api et al., 2015). As noted in the Criteria Document, the screening criteria applied are the same as those used in the EU for REACH (ECHA, 2012). For persistence, if the EPI Suite model BIOWIN 3 predicts a value < 2.2 and either BIOWIN 2 or BIOWIN 6 predicts a value < 0.5, then the material is considered potentially persistent. A material would be considered potentially bioaccumulative if the EPI Suite model BCFBAF predicts a fish BCF ≥ 2000 L/kg. Ecotoxicity is determined in the above screening-level risk assessment. If, based on these model outputs (Step 1), additional assessment is required, a WoE-based review is then performed (Step 2). This review considers

Exposure information and PEC calculation (following RIFM Environmental Framework: Salvito et al., 2002).

Exposure	Europe (EU)	North America (NA)
Log K_{ow} Used	4.07	4.07
Biodegradation Factor Used	0	0
Dilution Factor	3	3
Regional Volume of Use Tonnage Band	< 1	< 1
Risk Characterization: PEC/PNEC	< 1	< 1

Based on available data, the RQ for this material is < 1. No further assessment is necessary.

The RIFM PNEC is 0.00393 $\mu\text{g/L}$. The revised PEC/PNECs for EU and NA are: not applicable. The material was cleared at the screening-level and therefore does not present a risk to the aquatic environment at the current reported volumes of use.

Literature Search and Risk Assessment Completed On: 11/28/17.

11. Literature Search*

- **RIFM Database:** Target, Fragrance Structure Activity Group materials, other references, JECFA, CIR, SIDS
- **ECHA:** <http://echa.europa.eu/>
- **NTP:** <https://ntp.niehs.nih.gov/>
- OECD Toolbox

- **SciFinder:** <https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf>
- **PubMed:** <http://www.ncbi.nlm.nih.gov/pubmed>
- **TOXNET:** <http://toxnet.nlm.nih.gov/>
- **IARC:** <http://monographs.iarc.fr>
- **OECD SIDS:** <http://webnet.oecd.org/hpv/ui/Default.aspx>
- **EPA ACToR:** <https://actor.epa.gov/actor/home.xhtml>
- **US EPA HPVIS:** https://ofmpub.epa.gov/opphpv/public_search_publicdetails?submission_id=24959241&ShowComments=Yes&sqlstr=null&recordcount=0&User_title=DetailQuery%20Results&EndPointRpt=Y#submission
- **Japanese NITE:** <http://www.safe.nite.go.jp/english/db.html>
- **Japan Existing Chemical Data Base (JECDB):** http://dra4.nihs.go.jp/mhlw_data/jsp/SearchPageENG.jsp
- **Google:** <https://www.google.com>
- **ChemIDplus:** <https://chem.nlm.nih.gov/chemidplus/>

Search keywords: CAS number and/or material names.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fct.2019.110616>.



Appendix

Read-across Justification

Methods

The read-across analogs were identified following the strategy for structuring and reporting a read-across prediction of toxicity described in Schultz et al. (2015). The strategy is also consistent with the guidance provided by OECD within Integrated Approaches for Testing and Assessment (OECD, 2015) and the European Chemical Agency read-across assessment framework (ECHA, 2016).

- First, materials were clustered based on their structural similarity. Second, data availability and data quality on the selected cluster were examined. Third, appropriate read-across analogs from the cluster were confirmed by expert judgment.
- Tanimoto structure similarity scores were calculated using FCFC4 fingerprints (Rogers and Hahn, 2010).
- The physical–chemical properties of the target substance and the read-across analogs were calculated using EPI Suite (US EPA, 2012a).
- J_{\max} values were calculated using RIFM's skin absorption model (SAM). The parameters were calculated using the consensus model (Shen et al., 2014).
- DNA binding, mutagenicity, genotoxicity alerts and oncologic classification predictions were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- ER binding and repeat dose categorization were generated using OECD QSAR Toolbox v3.4 (OECD, 2012).
- Developmental toxicity was predicted using CAESAR v2.1.7 (Cassano et al., 2010), and skin sensitization was predicted using Toxtree 2.6.13.
- Protein binding was predicted using OECD QSAR Toolbox v3.4 (OECD, 2012).
- The major metabolites for the target and read-across analogs were determined and evaluated using OECD QSAR Toolbox v3.4 (OECD, 2012).

	Target Material	Read-across Material
Principal Name	Methyl 3,7-dimethyl-6-octenoate	<i>trans</i> -Methylgeranate
CAS No.	2270-60-2	1189-09-9
Structure		
Similarity (Tanimoto Score)		0.64
Read-across Endpoint		• Genotoxicity
Molecular Formula	$C_{11}H_{20}O_2$	$C_{11}H_{18}O_2$
Molecular Weight	184.28	182.26
Melting Point (°C, EPI Suite)	-18.46	-16.99
Boiling Point (°C, EPI Suite)	219.01	230.97
Vapor Pressure (Pa @ 25 °C, EPI Suite)	18.4	9.91
Log Kow (KOWWIN v1.68 in EPI Suite)	4.07	3.98
Water Solubility (mg/L, @ 25 °C, WSKOW v1.42 in EPI Suite)	17.56	21.24
J_{\max} ($\mu\text{g}/\text{cm}^2/\text{h}$, SAM)	30.993	38.586
Henry's Law ($\text{Pa}\cdot\text{m}^3/\text{mol}$, Bond Method, EPI Suite)	1.78E+002	9.72E-004
Genotoxicity		
DNA Binding (OASIS v1.4, QSAR Toolbox v3.4)	• No alert found	• No alert found

*Information sources outside of RIFM's database are noted as appropriate in the safety assessment. This is not an exhaustive list. The links listed above were active as of 08/02/2018.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome. RIFM staff are employees of the Research Institute for Fragrance Materials, Inc. (RIFM). The Expert Panel receives a small honorarium for time spent reviewing the subject work.

DNA Binding (OECD QSAR Toolbox v3.4)	● No alert found	● No alert found
Carcinogenicity (ISS)	● Non-toxicant (low reliability)	● Non-toxicant (low reliability)
DNA Binding (Ames, MN, CA, OASIS v1.1)	● No alert found	● No alert found
<i>In Vitro</i> Mutagenicity (Ames, ISS)	● No alert found	● No alert found
<i>In Vivo</i> Mutagenicity (Micronucleus, ISS)	● No alert found	● No alert found
Oncologic Classification	● Not classified	● Acrylate reactive functional groups
<i>Metabolism</i>		
Rat Liver S9 Metabolism Simulator and Structural Alerts for Metabolites (OECD QSAR Toolbox v3.4)	See Supplemental Data 1	See Supplemental Data 2

Summary

There are insufficient toxicity data on methyl 3,7-dimethyl-6-octenoate (CAS # 2270-60-2). Hence, *in silico* evaluation was conducted to determine read-across analogs for this material. Based on structural similarity, reactivity, metabolism, physical-chemical properties, and expert judgment, *trans*-methylgeranate (CAS # 1189-09-9) was identified as a read-across material with sufficient data for toxicological evaluation.

14. Conclusions

- *trans*-Methylgeranate (CAS # 1189-09-9) was used as a read-across analog for the target material methyl 3,7-dimethyl-6-octenoate (CAS # 2270-60-2) for the genotoxicity endpoint.
 - The target substance and the read-across analog are structurally similar and belong to the class of branched chain unsaturated esters.
 - The target substance and the read-across analog are methyl esters of similar branched chain acids.
 - The key structural difference between the target substance and the read-across analog is that the target substance has an additional α,β -unsaturation relative to the target material. This structural difference is toxicologically insignificant, making the read-across analog potentially more reactive.
 - Structural similarity between the target substance and the read-across analog is indicated by the Tanimoto score. The Tanimoto score reflects the similarity of these structures. Differences between the structures that affect the Tanimoto score are toxicologically insignificant.
 - The physical-chemical properties of the target substance and the read-across analog are sufficiently similar to enable comparison of their toxicological properties.
 - According to the OECD QSAR Toolbox v3.4, structural alerts for toxicological endpoints are consistent between the target substance and the read-across analog.
 - The read-across analog has an alert for oncologic classification. There is no such alert for the target substance. According to these predictions, the read-across analog is expected to be more reactive compared to the target substance. Data superseded predictions in this case.
 - The target substance and the read-across analog are expected to be metabolized similarly, as shown by the metabolism simulator.
 - The structural alerts for the endpoints evaluated are consistent between the metabolites of the read-across analog and the target material.

References

- Api, A.M., Belsito, D., Bruze, M., Cadby, P., Calow, P., Dagli, M.L., Dekant, W., Ellis, G., Fryer, A.D., Fukayama, M., Griem, P., Hickey, C., Kromidas, L., Lalko, J.F., Liebler, D.C., Miyachi, Y., Politano, V.T., Renskers, K., Ritacco, G., Salvito, D., Schultz, T.W., Sipes, I.G., Smith, B., Vitale, D., Wilcox, D.K., 2015. Criteria for the Research Institute for fragrance materials, Inc. (RIFM) safety evaluation process for fragrance ingredients. *Food Chem. Toxicol.* 82, S1–S19.
- Arctander, S., 1969. *Perfume and Flavor Chemicals (Aroma Chemicals)*, Vol. I and II. Published by the Author: Montclair, NJ (USA).
- Cassano, A., Manganaro, A., Martin, T., Young, D., Piclin, N., Pintore, M., Bigoni, D., Benfenati, E., 2010. CAESAR models for developmental toxicity. *Chem. Cent. J.* (4 Suppl. 1), S4.
- Carthew, P., Clapp, C., Gutsell, S., 2009. Exposure based waiving: the application of the toxicological threshold of concern (TTC) to inhalation exposure for aerosol ingredients in consumer products. *Food Chem. Toxicol.* 47 (6), 1287–1295.
- Comiskey, D., Api, A.M., Barratt, C., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Safford, B., Smith, B., Tozer, S., 2015. Novel database for exposure to fragrance ingredients in cosmetics and personal care products. *Regul. Toxicol. Pharmacol.* 72 (3), 660–672.
- Comiskey, D., Api, A.M., Barrett, C., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Rose, J., Safford, B., Smith, B., Tozer, S., 2017. Integrating habits and practices data for soaps, cosmetics and air care products into an existing aggregate exposure model. *Regul. Toxicol. Pharmacol.* 88, 144–156.
- ECHA, 2012. Guidance on information requirements and chemical safety assessment Chapter R.11: PBT Assessment, November 2012 v1.1. <http://echa.europa.eu/>.
- ECHA, 2016. European Chemical Agency Read-Across Assessment Framework. ECHA Read-across Assessment Framework. www.echa.europa.eu/documents/10162/13628/raaf_en.pdf.
- Henry, B., Foti, C., Alsante, K., 2009. Can light absorption and photostability data be used to assess the photosafety risks in patients for a new drug molecule? *J. Photochem. Photobiol. B Biol.* 96 (1), 57–62.
- IFRA (International Fragrance Association), 2015. Volume of Use Survey, February 2015.
- Kroes, R., Renwick, A.G., Feron, V., Galli, C.L., Gibney, M., Greim, H., Guy, R.H., Lhuguenot, J.C., van de Sandt, J.J.M., 2007. Application of the threshold of toxicological concern (TTC) to the safety evaluation of cosmetic ingredients. *Food Chem. Toxicol.* 45 (12), 2533–2562.
- Laufersweiler, M.C., Gadagbui, B., Baskerville-Abraham, I.M., Maier, A., Willis, A., et al., 2012. Correlation of chemical structure with reproductive and developmental toxicity as it relates to the use of the threshold of toxicological concern. *Regul. Toxicol. Pharmacol.* 62 (1), 160–182.
- OECD, 2012. The OECD QSAR Toolbox, V 3.4. <http://www.qsartoolbox.org/>.
- OECD, 2015. Guidance document on the reporting of integrated Approaches to testing and assessment. ENV/JM/HA(2015)7. Retrieved from. <http://www.oecd.org/>.
- RIFM (Research Institute for Fragrance Materials, Inc), 2013. Report on the Testing of Methyl 3,7-Dimethyl-6-Octenoate in the BlueScreen HC Assay (-/+ S9 Metabolic Activation). RIFM Report Number 66275. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2014a. *trans*-Methylgeranate (E-Methylgeranate): Bacterial Reverse Mutation Assay. RIFM Report Number 67353. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2014b. *trans*-Methylgeranate: in Vitro Micronucleus Assay in Human Peripheral Blood Lymphocytes. RIFM Report Number 67548. RIFM, Woodcliff Lake, NJ, USA.
- RIFM (Research Institute for Fragrance Materials, Inc), 2017. Exposure Survey 16 May 2017.
- Roberts, D.W., Api, A.M., Safford, R.J., Lalko, J.F., 2015. Principles for identification of high potency category chemicals for which the dermal sensitization threshold (DST) approach should not be applied. *Regul. Toxicol. Pharmacol.* 72 (3), 683–693.
- Rogers, D., Hahn, M., 2010. Extended-connectivity fingerprints. *J. Chem. Inf. Model.* 50 (5), 742–754.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Smith, B., Thomas, R., Tozer, S., 2015a. Use of an aggregate exposure model to estimate consumer exposure to fragrance ingredients in personal care and cosmetic products. *Regul. Toxicol. Pharmacol.* 72, 673–682.
- Safford, B., Api, A.M., Barratt, C., Comiskey, D., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Rose, J., Smith, B., Tozer, S., 2017. Application of the expanded Creme RIFM consumer exposure model to fragrance ingredients in cosmetic, personal care and air care products. *Regul. Toxicol. Pharmacol.* 86, 148–156.
- Safford, R.J., 2008. The dermal sensitisation threshold—A TTC approach for allergic contact dermatitis. *Regul. Toxicol. Pharmacol.* 51 (2), 195–200.
- Safford, R.J., Api, A.M., Roberts, D.W., Lalko, J.F., 2015b. Extension of the dermal sensitization threshold (DST) approach to incorporate chemicals classified as reactive. *Regul. Toxicol. Pharmacol.* 72 (3), 694–701.

- Safford, R.J., Aptula, A.O., Gilmour, N., 2011. Refinement of the dermal sensitisation threshold (DST) approach using a larger dataset and incorporating mechanistic chemistry domains. *Regul. Toxicol. Pharmacol.* 60 (2), 218–224.
- Salvito, D.T., Senna, R.J., Federle, T.W., 2002. A Framework for prioritizing fragrance materials for aquatic risk assessment. *Environ. Toxicol. Chem.* 21 (6), 1301–1308.
- Schultz, T.W., Amcoff, P., Berggren, E., Gautier, F., Klaric, M., Knight, D.J., Mahony, C., Schwarz, M., White, A., Cronin, M.T., 2015. A strategy for structuring and reporting a read-across prediction of toxicity. *Regul. Toxicol. Pharmacol.* 72 (3), 586–601.
- Shen, J., Kromidas, L., Schultz, T., Bhatia, S., 2014. An *in silico* skin absorption model for fragrance materials. *Food Chem. Toxicol.* 74, 164–176.
- US EPA, 2012a. Estimation Programs Interface Suite for Microsoft Windows, v4.0–v4.11. United States Environmental Protection Agency, Washington, DC, USA.
- US EPA, 2012b. The ECOSAR (ECOLOGICAL Structure Activity Relationship) Class Program for Microsoft Windows, v1.11. United States Environmental Protection Agency, Washington, DC, USA.