



Contents lists available at ScienceDirect

Food and Chemical Toxicology

journal homepage: www.elsevier.com/locate/foodchemtox

Short review



RIFM fragrance ingredient safety assessment, benzaldehyde glyceryl acetal, CAS Registry Number 1319-88-6

A.M. Api^a, D. Belsito^b, S. Biserta^a, D. Botelho^a, M. Bruze^c, G.A. Burton Jr.^d, J. Buschmann^e, M. A. Cancellieri^a, M.L. Dagli^f, M. Date^a, W. Dekant^g, C. Deodhar^a, A.D. Fryer^h, S. Gadhia^a, L. Jones^a, K. Joshi^a, M. Kumar^a, A. Lapczynski^a, M. Lavelle^a, I. Lee^a, D.C. Lieblerⁱ, H. Moustakas^a, M. Na^a, T.M. Penning^j, G. Ritacco^a, J. Romine^a, N. Sadekar^a, T.W. Schultz^k, D. Selechnik^a, F. Siddiqi^a, I.G. Sipes^l, G. Sullivan^{a,*}, Y. Thakkar^a, Y. Tokura^m

^a Research Institute for Fragrance Materials, Inc., 50 Tice Boulevard, Woodcliff Lake, NJ, 07677, USA^b Columbia University Medical Center, Department of Dermatology, 161 Fort Washington Ave., New York, NY, 10032, USA^c Malmo University Hospital, Department of Occupational & Environmental Dermatology, Sodra Forstadsgatan 101, Entrance 47, Malmo, SE, 20502, Sweden^d School of Natural Resources & Environment, University of Michigan, Dana Building G110, 440 Church St., Ann Arbor, MI, 48109, USA^e Fraunhofer Institute for Toxicology and Experimental Medicine, Nikolai-Fuchs-Strasse 1, 30625, Hannover, Germany^f University of Sao Paulo, School of Veterinary Medicine and Animal Science, Department of Pathology, Av. Prof. dr. Orlando Marques de Paiva, 87, Sao Paulo, CEP 05508-900, Brazil^g University of Wuerzburg, Department of Toxicology, Versbacher Str. 9, 97078, Würzburg, Germany^h Oregon Health Science University, 3181 SW Sam Jackson Park Rd., Portland, OR, 97239, USAⁱ Vanderbilt University School of Medicine, Department of Biochemistry, Center in Molecular Toxicology, 638 Robinson Research Building, 2200 Pierce Avenue, Nashville, TN, 37232-0146, USA^j University of Pennsylvania, Perelman School of Medicine, Center of Excellence in Environmental Toxicology, 1316 Biomedical Research Building (BRB) II/III, 421 Curie Boulevard, Philadelphia, PA, 19104-3083, USA^k The University of Tennessee, College of Veterinary Medicine, Department of Comparative Medicine, 2407 River Dr., Knoxville, TN, 37996-4500, USA^l Department of Pharmacology, University of Arizona, College of Medicine, 1501 North Campbell Avenue, P.O. Box 245050, Tucson, AZ, 85724-5050, USA^m The Journal of Dermatological Science (JDS), Department of Dermatology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, 431-3192, Japan

ARTICLE INFO

Handling editor: Aristides Tsatsakis

Keywords:

Genotoxicity
Repeated dose, developmental, and reproductive toxicity
Skin sensitization
Phototoxicity/photoallergenicity
Local respiratory toxicity
Environmental safety

(continued)

Version: 110420. This version replaces any previous versions.

Name: Benzaldehyde glyceryl acetal CAS Registry Number: 1319-88-6

(continued on next column)

(continued on next page)

* Corresponding author.


E-mail address: gsullivan@rifm.org (G. Sullivan).<https://doi.org/10.1016/j.fct.2021.112173>

Received 5 November 2020; Received in revised form 3 March 2021; Accepted 30 March 2021

Available online 14 April 2021

0278-6915/© 2021 Elsevier Ltd. All rights reserved.

(continued)

**Abbreviation/Definition List:**

2-Box Model - A RIFM, Inc. proprietary *in silico* tool used to calculate fragrance air exposure concentration

AF - Assessment Factor

BCF - Bioconcentration Factor

Crema RIFM Model - The Crema RIFM Model uses probabilistic (Monte Carlo) simulations to allow full distributions of data sets, providing a more realistic estimate of aggregate exposure to individuals across a population (Comiskey et al., 2015, 2017; bib.Comiskey_et_al_2015; bib.Comiskey_et_al_2015; Safford et al., 2015a, 2017; bib.Safford_et_al_2015; bib.Safford_et_al_2017; bib.Comiskey_et_al_2017) compared to a deterministic aggregate approach

DEREK - Derek Nexus is an *in silico* tool used to identify structural alerts

DRF - Dose Range Finding

DST - Dermal Sensitization Threshold

ECHA - European Chemicals Agency

ECOSAR - Ecological Structure-Activity Relationships Predictive Model

EU - Europe/European Union

GLP - Good Laboratory Practice

IFRA - The International Fragrance Association

LOEL - Lowest Observable Effect Level

MOE - Margin of Exposure

MPPD - Multiple-Path Particle Dosimetry. An *in silico* model for inhaled vapors used to simulate fragrance lung deposition

NA - North America

NESIL - No Expected Sensitization Induction Level

NOAEC - No Observed Adverse Effect Concentration

NOAEL - No Observed Adverse Effect Level

NOEC - No Observed Effect Concentration

NOEL - No Observed Effect Level

OECD - Organisation for Economic Co-operation and Development

OECD TG - Organisation for Economic Co-operation and Development Testing Guidelines

PBT - Persistent, Bioaccumulative, and Toxic

PEC/PNEC - Predicted Environmental Concentration/Predicted No Effect Concentration

Perfumery - In this safety assessment, perfumery refers to fragrances made by a perfumer used in consumer products only. The exposures reported in the safety assessment include consumer product use but do not include occupational exposures.

QRA - Quantitative Risk Assessment

QSAR - Quantitative Structure-Activity Relationship

REACH - Registration, Evaluation, Authorisation, and Restriction of Chemicals

RfD - Reference Dose

RIFM - Research Institute for Fragrance Materials

RQ - Risk Quotient

Statistically Significant - Statistically significant difference in reported results as compared to controls with a $p < 0.05$ using appropriate statistical test

TTC - Threshold of Toxicological Concern

UV/Vis spectra - Ultraviolet/Visible spectra

VCF - Volatile Compounds in Food

VoU - Volume of Use

vPvB - (very) Persistent, (very) Bioaccumulative

WoE - Weight of Evidence

The Expert Panel for Fragrance Safety* concludes that this material is safe as described in this safety assessment.

This safety assessment is based on the RIFM Criteria Document (Api, 2015), which should be referred to for clarifications.

Each endpoint discussed in this safety assessment includes the relevant data that were available at the time of writing (version number in the top box is indicative of the date of approval based on a 2-digit month/day/year), both in the RIFM Database (consisting of publicly available and proprietary data) and through publicly available information sources (e.g., SciFinder and PubMed). Studies selected for this safety assessment were based on appropriate test criteria, such as acceptable

(continued on next column)

(continued)

guidelines, sample size, study duration, route of exposure, relevant animal species, most relevant testing endpoints, etc. A key study for each endpoint was selected based on the most conservative endpoint value (e.g., PNEC, NOAEL, LOEL, and NESIL).

*The Expert Panel for Fragrance Safety is an independent body that selects its own members and establishes its own operating procedures. The Expert Panel is comprised of internationally known scientists that provide RIFM with guidance relevant to human health and environmental protection.

Summary: The existing information supports the use of this material as described in this safety assessment.

Benzaldehyde glyceryl acetal was evaluated for genotoxicity, repeated dose toxicity, reproductive toxicity, local respiratory toxicity, phototoxicity/photoallergenicity, skin sensitization, and environmental safety. Data show that benzaldehyde glyceryl acetal is not genotoxic. The repeated dose, reproductive, and local respiratory toxicity endpoints were evaluated using the Threshold of Toxicological Concern (TTC) for a Cramer Class II material, and the exposure to benzaldehyde glyceryl acetal is below the TTC (0.009 mg/kg/day, 0.009 mg/kg/day, and 0.47 mg/day, respectively). The skin sensitization endpoint was completed using the Dermal Sensitization Threshold (DST) for non-reactive materials (900 $\mu\text{g}/\text{cm}^2$); exposure is below the DST. The phototoxicity/photoallergenicity endpoints were evaluated based on ultraviolet/visible (UV/Vis) spectra; benzaldehyde glyceryl acetal is not expected to be phototoxic/photoallergenic. The environmental endpoints were evaluated; benzaldehyde glyceryl acetal was found not to be Persistent, Bioaccumulative, and Toxic (PBT) as per the International Fragrance Association (IFRA) Environmental Standards, and its risk quotients, based on its current volume of use in Europe and North America (i.e., Predicted Environmental Concentration/Predicted No Effect Concentration [PEC/PNEC]), are <1 .

Human Health Safety Assessment

Genotoxicity: Not genotoxic. (RIFM, 2016b)

Repeated Dose Toxicity: No NOAEL available. Exposure is below TTC.

Reproductive Toxicity: No NOAEL available. Exposure is below TTC.

Skin Sensitization: No safety concerns at current, declared use levels; Exposure is below the DST.

Phototoxicity/Photoallergenicity: Not expected to be phototoxic/photoallergenic. (UV Spectra; RIFM Database)

Local Respiratory Toxicity: No NOAEC available. Exposure is below the TTC.

Environmental Safety Assessment**Hazard Assessment:**

Persistence:
Screening-level: 2.97 (BIOWIN 3) (EPI Suite v4.11; US EPA, 2012a)

Bioaccumulation:
Screening-level: 3.16 L/kg (EPI Suite v4.11; US EPA, 2012a)

Ecotoxicity:
Screening-level: Fish LC50: 7623 mg/L (RIFM Framework; Salvito, 2002)

Conclusion: Not PBT or vPvB as per IFRA Environmental Standards

Risk Assessment:

Screening-level: PEC/PNEC (North America and Europe) <1 (RIFM Framework; Salvito, 2002)

Critical Ecotoxicity Endpoint: Fish LC50: 7623 mg/L (RIFM Framework; Salvito, 2002)

RIFM PNEC is: 7.623 $\mu\text{g}/\text{L}$

- Revised PEC/PNECs (2015 IFRA VoU): North America and Europe: not applicable; cleared at screening-level

1. Identification

- 1. Chemical Name:** Benzaldehyde glyceryl acetal
- 2. CAS Registry Number:** 1319-88-6
- 3. Synonyms:** Benzaldehyde, cyclic acetal with glycerol; Benzalglycerin; 1,3-Dioxolane-4-methanol, 2-phenyl-; 4-Hydroxymethyl-2-phenyl-m-dioxolane; 5-Hydroxy-2-phenyl-1,3-dioxane; 2-Phenyl-1,3-dioxan-5-ol; 2-Phenyl-m-dioxan-5-ol; Benzaldehyde glyceryl acetal
- 4. Molecular Formula:** $\text{C}_{10}\text{H}_{12}\text{O}_3$
- 5. Molecular Weight:** 180.2
- 6. RIFM Number:** 495
- 7. Stereochemistry:** Isomer not specified. Two chiral centers present and a total of 4 enantiomers possible.

2. Physical data

- Boiling Point:** 280 °C (Fragrance Materials Association [FMA]), 298.18 °C (EPI Suite)
- Flash Point:** >93 °C (Globally Harmonized System), >200 °F; CC (FMA)
- Log K_{ow}:** 0.28 (EPI Suite)
- Melting Point:** 66.63 °C (EPI Suite)
- Water Solubility:** 102,500 mg/L (EPI Suite)
- Specific Gravity:** 1.190 (FMA)
- Vapor Pressure:** 0.0000297 mm Hg at 20 °C (EPI Suite v4.0), 0.01 mm Hg at 20 °C (FMA), 6.17e-005 mm Hg at 25 °C (EPI Suite)
- UV Spectra:** No significant absorbance between 290 and 700 nm; molar absorption coefficient is below the benchmark (1000 L mol⁻¹ · cm⁻¹)
- Appearance/Organoleptic:** A colorless viscous liquid with a very faint odor

3. Volume of use (worldwide band)

- 0.1–1 metric ton per year (IFRA, 2015)

4. Exposure to fragrance ingredient (Creme RIFM Aggregate Exposure Model v3.0)

- 95th Percentile Concentration in Scented Candles:** 1.5% (RIFM, 2019)

No reported use in hydroalcoholics

- Inhalation Exposure*:** 0.00019 mg/kg/day or 0.014 mg/day (RIFM, 2019)
- Total Systemic Exposure**:** 0.00021 mg/kg/day (RIFM, 2019)

*95th percentile calculated exposure derived from concentration survey data in the Creme RIFM Aggregate Exposure Model (Comiskey, 2015, 2017bib_Comiskey_et_al_2015bib_Comiskey_et_al_2015; Safford, 2015a, 2017bib_Safford_et_al_2015bib_Safford_et_al_2017bib_Comiskey_et_al_2017).

**95th percentile calculated exposure; assumes 100% absorption unless modified by dermal absorption data as reported in Section V. It is derived from concentration survey data in the Creme RIFM Aggregate Exposure Model and includes exposure via dermal, oral, and inhalation routes whenever the fragrance ingredient is used in products that include these routes of exposure (Comiskey, 2015, 2017bib_Comiskey_et_al_2015bib_Comiskey_et_al_2015; Safford, 2015a, 2017bib_Safford_et_al_2015bib_Safford_et_al_2017bib_Comiskey_et_al_2017).

5. Derivation of systemic absorption

- Dermal:** Assumed 100%
- Oral:** Assumed 100%
- Inhalation:** Assumed 100%

6. Computational toxicology evaluation

- Cramer Classification:** Class II, Intermediate (Expert Judgment)

Expert Judgment	Toxtree v3.1	OECD QSAR Toolbox v3.2
II*	III	III

*Due to potential discrepancies with the current *in silico* tools (Bhatia et al., 2015), the Cramer Class of the target material was determined using expert judgment based on the Cramer decision tree (Cramer et al., 1978). See the Appendix below for further details.

2. Analogs Selected:

- Genotoxicity:** None
 - Repeated Dose Toxicity:** None
 - Reproductive Toxicity:** None
 - Skin Sensitization:** None
 - Phototoxicity/Photoallergenicity:** None
 - Local Respiratory Toxicity:** None
 - Environmental Toxicity:** None
3. **Read-across Justification:** None

7. Metabolism

No relevant data available for inclusion in this safety assessment.
Additional References: None.

8. Natural occurrence

Benzaldehyde glyceryl acetal is not reported to occur in foods by the VCF*.

*VCF (Volatile Compounds in Food): Database/Nijssen, L.M.; Ingen-Visscher, C.A. van; Donders, J.J.H. (eds). – Version 15.1 – Zeist (The Netherlands): TNO Triskelion, 1963–2014. A continually updated database containing information on published volatile compounds that have been found in natural (processed) food products. Includes FEMA GRAS and EU-Flavis data.

9. REACH dossier

Pre-registered for 2010; no dossier available as of 11/04/20.

10. Conclusion

The existing information supports the use of this material as described in this safety assessment.

11. Summary

11.1. Human health endpoint summaries

11.1.1. Genotoxicity

Based on the current existing data, benzaldehyde glyceryl acetal does not present a concern for genotoxicity.

11.1.1.1. Risk assessment. Benzaldehyde glyceryl acetal was assessed in the BlueScreen assay and found positive for cytotoxicity (positive: <80% relative cell density) and negative for genotoxicity without metabolic activation, and negative for both cytotoxicity (positive: <80% relative cell density) and genotoxicity with metabolic activation (RIFM, 2013). BlueScreen is a human cell-based assay for measuring the genotoxicity and cytotoxicity of chemical compounds and mixtures. Additional assays were considered to fully assess the potential mutagenic or clastogenic effects of the target material.

The mutagenic activity of benzaldehyde glyceryl acetal has been evaluated in a bacterial reverse mutation assay conducted in compliance with GLP regulations and in accordance with OECD TG 471 using the standard plate incorporation method. *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and *Escherichia coli* strain WP2uvrA were treated with benzaldehyde glyceryl acetal in dimethyl sulfoxide (DMSO) at concentrations up to 5000 µg/plate. No increases in the mean number of revertant colonies were observed at any tested concentration in the presence or absence of S9 (RIFM, 2016a). Under the conditions of the study, benzaldehyde glyceryl acetal was not mutagenic in the Ames test.

The clastogenic activity of benzaldehyde glyceryl acetal was evaluated in an *in vitro* micronucleus test conducted in compliance with GLP

regulations and in accordance with OECD TG 487. Human peripheral blood lymphocytes were treated with benzaldehyde glyceryl acetal in DMSO, and micronuclei analysis was conducted at concentrations up to 10,000 μM in the presence and absence of metabolic activation. Benzaldehyde glyceryl acetal did not induce binucleated cells with micronuclei when tested up to cytotoxic levels or the maximum concentration in either the presence or absence of an S9 activation system (RIFM, 2016b). Under the conditions of the study, benzaldehyde glyceryl acetal was considered to be non-clastogenic in the *in vitro* micronucleus test.

Based on the data available, benzaldehyde glyceryl acetal does not present a concern for genotoxic potential.

Additional References: None.

Literature Search and Risk Assessment Completed On: 02/19/20.

11.1.2. Repeated dose toxicity

There are no repeated dose toxicity data on benzaldehyde glyceryl acetal or any read-across materials. The total systemic exposure to benzaldehyde glyceryl acetal is below the TTC for the repeated dose toxicity endpoint of a Cramer Class II material at the current level of use.

11.1.2.1. Risk assessment. There are no repeated dose toxicity data on benzaldehyde glyceryl acetal or any read-across materials that can be used to support the repeated dose toxicity endpoint. The total systemic exposure (0.21 $\mu\text{g}/\text{kg}/\text{day}$) is below the TTC for benzaldehyde glyceryl acetal (9 $\mu\text{g}/\text{kg}/\text{day}$; Kroes, 2007).

Additional References: None

Literature Search and Risk Assessment Completed On: 02/26/20.

11.1.3. Reproductive toxicity

There are insufficient reproductive toxicity data on benzaldehyde glyceryl acetal or any read-across materials. The total systemic exposure to benzaldehyde glyceryl acetal is below the TTC for the reproductive toxicity endpoint of a Cramer Class II material at the current level of use.

11.1.3.1. Risk assessment. There are no reproductive toxicity data on benzaldehyde glyceryl acetal or any read-across materials that can be used to support the reproductive toxicity endpoint. The total systemic exposure (0.21 $\mu\text{g}/\text{kg}/\text{day}$) is below the TTC for benzaldehyde glyceryl acetal (9 $\mu\text{g}/\text{kg}/\text{day}$; Kroes, 2007; Laufersweiler, 2012).

Additional References: None.

Literature Search and Risk Assessment Completed On: 03/19/20.

11.1.4. Skin sensitization

Based on existing data and the application of DST, benzaldehyde glyceryl acetal does not present a safety concern for skin sensitization under the current, declared levels of use.

11.1.4.1. Risk assessment. Limited skin sensitization studies are available for benzaldehyde glyceryl acetal. The chemical structure of this material indicates that it would not be expected to react with skin proteins directly, while its metabolite is expected to be reactive (Roberts, 2007; Toxtree v3.1.0; OECD Toolbox v4.2). In a human maximization test, no skin sensitization reactions were observed with 4% and 8% benzaldehyde glyceryl acetal (RIFM, 1980; RIFM, 1974). Due to the limited data, the reported exposure was benchmarked utilizing the non-reactive DST of 900 $\mu\text{g}/\text{cm}^2$ (Safford, 2008, 2011, 2015bib_Safford_2008bib_Safford_et_al_2011; Roberts, 2015bib_Safford_et_al_2015b). The current exposure from the 95th percentile concentration is below the DST for non-reactive materials when evaluated in all QRA categories. Table 1 provides the maximum acceptable concentrations for benzaldehyde glyceryl acetal that present no

Table 1

Maximum acceptable concentrations for benzaldehyde glyceryl acetal that present no appreciable risk for skin sensitization based on non-reactive DST.

IFRA Category ^a	Description of Product Type	Maximum Acceptable Concentrations in Finished Products Based on Non-reactive DST	Reported 95th Percentile Use Concentrations in Finished Products
1	Products applied to the lips	0.069%	NRU ^b
2	Products applied to the axillae	0.021%	NRU ^b
3	Products applied to the face using fingertips	0.41%	NRU ^b
4	Fine fragrance products	0.39%	NRU ^b
5	Products applied to the face and body using the hands (palms), primarily leave-on	0.10%	NRU ^b
6	Products with oral and lip exposure	0.23%	NRU ^b
7	Products applied to the hair with some hand contact	0.79%	NRU ^b
8	Products with significant anogenital exposure	0.041%	No Data ^c
9	Products with body and hand exposure, primarily rinse-off	0.75%	0.003%
10	Household care products with mostly hand contact	2.7%	NRU ^b
11	Products with intended skin contact but minimal transfer of fragrance to skin from inert substrate	1.5%	No Data ^c
12	Products not intended for direct skin contact, minimal or insignificant transfer to skin	No Restriction	1.5%

^a For a description of the categories, refer to the IFRA/RIFM Information Booklet.

^b No reported use.

^c Fragrance exposure from these products is very low. These products are not currently in the Creme RIFM Aggregate Exposure Model.

appreciable risk for skin sensitization based on the non-reactive DST. These levels represent maximum acceptable concentrations based on the DST approach. However, additional studies may show it could be used at higher levels.

Additional References: None.

Literature Search and Risk Assessment Completed On: 04/07/20.

11.1.5. Phototoxicity/photoallergenicity

Based on the available UV/Vis spectra, benzaldehyde glyceryl acetal would not be expected to present a concern for phototoxicity or photoallergenicity.

11.1.5.1. Risk assessment. There are no phototoxicity studies available for benzaldehyde glyceryl acetal in experimental models. UV/Vis absorption spectra indicate no significant absorption between 290 and 700 nm. The corresponding molar absorption coefficient is well below the benchmark of concern for phototoxicity and photoallergenicity (Henry, 2009). Based on the lack of absorbance, benzaldehyde glyceryl

acetal does not present a concern for phototoxicity or photoallergenicity.

11.1.5.2. UV spectra analysis. UV/Vis absorption spectra (OECD TG 101) were obtained. The spectra indicate no significant absorbance in the range of 290–700 nm. The molar absorption coefficient is below the benchmark of concern for phototoxic effects, $1000 \text{ L mol}^{-1} \cdot \text{cm}^{-1}$ (Henry, 2009).

Additional References: None.

Literature Search and Risk Assessment Completed On: 03/20/20.

11.1.6. Local respiratory toxicity

The margin of exposure could not be calculated due to a lack of appropriate data. The exposure level for benzaldehyde glyceryl acetal is below the Cramer Class III* TTC value for inhalation exposure local effects.

11.1.6.1. Risk assessment. There are no inhalation data available on benzaldehyde glyceryl acetal. Based on the Creme RIFM Model, the inhalation exposure is 0.014 mg/day. This exposure is 33.6 times lower than the Cramer Class III* TTC value of 0.47 mg/day (based on human lung weight of 650 g; Carthew, 2009); therefore, the exposure at the current level of use is deemed safe.

*As per Carthew et al. (2009), Cramer Class II materials default to Cramer Class III for the local respiratory toxicity endpoint.

Additional References: None.

Literature Search and Risk Assessment Completed On: 04/01/20.

11.2. Environmental endpoint summary

11.2.1. Screening-level assessment

A screening-level risk assessment of benzaldehyde glyceryl acetal was performed following the RIFM Environmental Framework (Salvito, 2002), which provides 3 tiered levels of screening for aquatic risk. In Tier 1, only the material's regional VoU, its log K_{OW} , and its molecular weight are needed to estimate a conservative risk quotient (RQ), expressed as the ratio Predicted Environmental Concentration/Predicted No Effect Concentration (PEC/PNEC). A general QSAR with a high uncertainty factor applied is used to predict fish toxicity, as discussed in Salvito et al. (2002). In Tier 2, the RQ is refined by applying a lower uncertainty factor to the PNEC using the ECOSAR model (US EPA, 2012b), which provides chemical class-specific ecotoxicity estimates. Finally, if necessary, Tier 3 is conducted using measured biodegradation and ecotoxicity data to refine the RQ, thus allowing for lower PNEC uncertainty factors. The data for calculating the PEC and PNEC for this safety assessment are provided in the table below. For the PEC, the range from the most recent IFRA Volume of Use Survey is reviewed. The PEC is then calculated using the actual regional tonnage, not the extremes of the range. Following the RIFM Environmental Framework, benzaldehyde glyceryl acetal was identified as a fragrance material with no potential to present a possible risk to the aquatic environment (i.e., its screening-level PEC/PNEC < 1).

A screening-level hazard assessment using EPI Suite v4.11 (US EPA, 2012a) did not identify benzaldehyde glyceryl acetal as possibly persistent or bioaccumulative based on its structure and physical-chemical properties. This screening-level hazard assessment considers the potential for a material to be persistent and bioaccumulative and toxic, or very persistent and very bioaccumulative as defined in the Criteria Document (Api, 2015). As noted in the Criteria Document, the screening criteria applied are the same as those used in the EU for REACH (ECHA, 2012). For persistence, if the EPI Suite model BIOWIN 3 predicts a value < 2.2 and either BIOWIN 2 or BIOWIN 6 predicts a value < 0.5, then the material is considered potentially persistent. A

material would be considered potentially bioaccumulative if the EPI Suite model BCFBAF predicts a fish BCF $\geq 2000 \text{ L/kg}$. Ecotoxicity is determined in the above screening-level risk assessment. If, based on these model outputs (Step 1), additional assessment is required, a WoE-based review is then performed (Step 2). This review considers available data on the material's physical-chemical properties, environmental fate (e.g., OECD Guideline biodegradation studies or die-away studies), fish bioaccumulation, and higher-tier model outputs (e.g., US EPA's BIOWIN and BCFBAF found in EPI Suite v4.11).

11.2.2. Risk assessment

Based on the current Volume of Use (2015), benzaldehyde glyceryl acetal presents no risk to the aquatic compartment in the screening-level assessment.

11.2.2.1. Key studies

11.2.2.1.1. Biodegradation. No data available.

11.2.2.1.2. Ecotoxicity. No data available.

11.2.2.1.3. Other available data. Benzaldehyde glyceryl acetal has been pre-registered for REACH with no additional information available at this time.

11.2.3. Risk assessment refinement

Ecotoxicological data and PNEC derivation (all endpoints reported in mg/L; PNECs in $\mu\text{g/L}$).

Endpoints used to calculate PNEC are underlined.

Exposure information and PEC calculation (following RIFM Environmental Framework: Salvito, 2002).

Exposure	Europe (EU)	North America (NA)
Log K_{OW} Used	0.28	0.28
Biodegradation Factor Used	0	0
Dilution Factor	3	3
Regional Volume of Use Tonnage Band	<1	<1
Risk Characterization: PEC/PNEC	<1	<1

Based on available data, the RQ for this material is < 1. No further assessment is necessary.

The RIFM PNEC is 7.623 $\mu\text{g/L}$. The revised PEC/PNECs for EU and NA are not applicable. The material was cleared at the screening-level; therefore, it does not present a risk to the aquatic environment at the current reported volumes of use.

Literature Search and Risk Assessment Completed On: 03/16/20.

12. Literature Search*

- **RIFM Database:** Target, Fragrance Structure-Activity Group materials, other references, JECFA, CIR, SIDS
- **ECHA:** <https://echa.europa.eu/>
- **NTP:** <https://ntp.niehs.nih.gov/>
- **OECD Toolbox:** <https://www.oecd.org/chemicalsafety/risk-assessment/oecd-qsar-toolbox.htm>
- **SciFinder:** <https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf>
- **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed>
- **National Library of Medicine's Toxicology Information Services:** <https://toxnet.nlm.nih.gov/>
- **IARC:** <https://monographs.iarc.fr>
- **OECD SIDS:** <https://hpvchemicals.oecd.org/ui/Default.aspx>
- **EPA ACToR:** <https://actor.epa.gov/actor/home.xhtml>
- **US EPA HPVIS:** https://ofmpub.epa.gov/opppvp/public_search_publicdetails?submission_id=24959241&ShowComments=Yes&sqlstr=null&recordcount=0&User_title=DetailQuery%20Results&EndPointRpt=Y#submission

		(mg/L)	(mg/L)			
RIFM Framework Screening-level (Tier 1)	7623			1000000	7.623	

- **Japanese NITE:** https://www.nite.go.jp/en/chem/chrip/chrip_search/systemTop
- **Japan Existing Chemical Data Base (JECDB):** http://dra4.nihs.go.jp/mhlw_data/jsp/SearchPageENG.jsp
- **Google:** <https://www.google.com>
- **ChemIDplus:** <https://chem.nlm.nih.gov/chemidplus/>

Search keywords: CAS number and/or material names.

*Information sources outside of RIFM's database are noted as appropriate in the safety assessment. This is not an exhaustive list. The links listed above were active as of 09/30/20.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome. RIFM staff are employees of the Research Institute for Fragrance Materials, Inc. (RIFM). The Expert Panel receives a small honorarium for time spent reviewing the subject work.

Appendix

Explanation of Cramer Classification

Due to potential discrepancies with the current *in silico* tools (Bhatia et al., 2015), the Cramer class of the target material was determined using expert judgment based on the Cramer decision tree (Cramer et al., 1978).

- Q1 A normal constituent of the body? No.
 Q2 Contains functional groups associated with enhanced toxicity? No.
 Q3 Contains elements other than C, H, O, N, and divalent S? No.
 Q5 Simply branched aliphatic hydrocarbon or a common carbohydrate? No.
 Q6 Benzene derivative with certain substituents? No.
 Q7 Heterocyclic? No.
 Q8 Lactone or cyclic diester? No.
 Q10 3-Membered heterocycles? No.
 Q11 Has a heterocyclic ring with complex substituents? No.
 Q12 Heteroaromatic? No.
 Q22 A common component of food? Yes. Class III (Class High)

References

Api, A.M., Belsito, D., Bruze, M., Cadby, P., Calow, P., Dagli, M.L., Dekant, W., Ellis, G., Fryer, A.D., Fukayama, M., Griem, P., Hickey, C., Kromidas, L., Lalko, J.F., Liebler, D.C., Miyachi, Y., Politano, V.T., Renskers, K., Ritacco, G., Salvito, D., Schultz, T.W., Sipes, I.G., Smith, B., Vitale, D., Wilcox, D.K., 2015. Criteria for the Research Institute for fragrance materials, Inc. (RIFM) safety evaluation process for fragrance ingredients. *Food Chem. Toxicol.* 82, S1–S19.

Bhatia, S., Schultz, T., Roberts, D., Shen, J., Kromidas, L., Api, A.M., 2015. Comparison of cramer classification between toxtree, the OECD QSAR Toolbox and expert judgment. *Regul. Toxicol. Pharmacol.* 71 (1), 52–62.

Carthew, P., Clapp, C., Gutsell, S., 2009. Exposure based waiving: the application of the toxicological threshold of concern (TTC) to inhalation exposure for aerosol ingredients in consumer products. *Food Chem. Toxicol.* 47 (6), 1287–1295.

Comiskey, D., Api, A.M., Barratt, C., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S.H., Safford, B., Smith, B., Tozer, S., 2015. Novel database for exposure to fragrance ingredients in cosmetics and personal care products. *Regul. Toxicol. Pharmacol.* 72 (3), 660–672.

Comiskey, D., Api, A.M., Barrett, C., Ellis, G., McNamara, C., O'Mahony, C., Robison, S. H., Rose, J., Safford, B., Smith, B., Tozer, S., 2017. Integrating habits and practices data for soaps, cosmetics and air care products into an existing aggregate exposure model. *Regul. Toxicol. Pharmacol.* 88, 144–156.

Cramer, G.M., Ford, R.A., Hall, R.L., 1978. Estimation of toxic hazard—a decision tree approach. *Food Chem. Toxicol.* 16 (3), 255–276.

Echa, 2012. Guidance on Information Requirements and Chemical Safety Assessment Chapter R.11: PBT Assessment, November 2012 v1.1. <http://echa.europa.eu/>.

Henry, B., Foti, C., Alsante, K., 2009. Can light absorption and photostability data be used to assess the photosafety risks in patients for a new drug molecule? *J. Photochem. Photobiol. B Biol.* 96 (1), 57–62.

Iifa (International Fragrance Association), 2015. Volume of Use Survey. February 2015.

Kroes, R., Renwick, A.G., Feron, V., Galli, C.L., Gibben, M., Greim, H., Guy, R.H., Lhuguenot, J.C., van de Sandt, J.J.M., 2007. Application of the threshold of toxicological concern (TTC) to the safety evaluation of cosmetic ingredients. *Food Chem. Toxicol.* 45 (12), 2533–2562.

Laufersweiler, M.C., Gadagbui, B., Baskerville-Abraham, I.M., Maier, A., Willis, A., et al., 2012. Correlation of chemical structure with reproductive and developmental toxicity as it relates to the use of the threshold of toxicological concern. *Regul. Toxicol. Pharmacol.* 62 (1), 160–182.

Rifm (Research Institute for Fragrance Materials, Inc), 1974. Report on Human Maximization Studies. Report to RIFM. RIFM Report Number 1801. RIFM, Woodcliff Lake, NJ, USA.

Rifm (Research Institute for Fragrance Materials, Inc), 1980. The appraisal of the contact-sensitizing potential of fragrance materials in humans. Report to RIFM. RIFM Report Number 1791. RIFM, Woodcliff Lake, NJ, USA.

Rifm (Research Institute for Fragrance Materials, Inc), 2013. Report on the Testing of Benzaldehyde Glyceryl Acetal in the BlueScreen HC Assay (-/+ S9 Metabolic Activation). RIFM Report Number 66187. RIFM, Woodcliff Lake, NJ, USA.

Rifm (Research Institute for Fragrance Materials, Inc), 2016a. Benzaldehyde Glyceryl Acetal: Genetic Toxicity Evaluation Using a Bacterial Reverse Mutation Test in Salmonella typhimurium TA1535, TA1537, TA98 and TA100, and Escherichia coli WP2 uvrA/pKM101. RIFM Report Number 70833. RIFM, Woodcliff Lake, NJ, USA.

Rifm (Research Institute for Fragrance Materials, Inc), 2016b. Benzaldehyde Glyceryl Acetal: Genetic Toxicity Evaluation Using a Micronucleus Test in Human Lymphocyte Cells. RIFM Report Number 71369. RIFM, Woodcliff Lake, NJ, USA.

Rifm (Research Institute for Fragrance Materials, Inc), 2019. Exposure Survey 25, October 2019.

Roberts, D.W., Api, A.M., Safford, R.J., Lalko, J.F., 2015. Principles for identification of high potency category chemicals for which the dermal sensitization threshold (DST) approach should not be applied. *Regul. Toxicol. Pharmacol.* 72 (3), 683–693.

Roberts, D.W., Patlewicz, G., Kern, P.S., Gerberick, F., Kimber, I., Dearman, R.J., Ryan, C. A., Basketter, D.A., Aptula, A.O., 2007. Mechanistic applicability domain classification of a local lymph node assay dataset for skin sensitization. *Chem. Res. Toxicol.* 20 (7), 1019–1030.

Safford, B., Api, A.M., Barratt, C., Comiskey, D., Daly, E.J., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Smith, B., Thomas, R., Tozer, S., 2015b. Use of an aggregate exposure model to estimate consumer exposure to fragrance ingredients in personal care and cosmetic products. *Regul. Toxicol. Pharmacol.* 72, 673–682.

Safford, B., Api, A.M., Barratt, C., Comiskey, D., Ellis, G., McNamara, C., O'Mahony, C., Robison, S., Rose, J., Smith, B., Tozer, S., 2017. Application of the expanded Creme RIFM consumer exposure model to fragrance ingredients in cosmetic, personal care and air care products. *Regul. Toxicol. Pharmacol.* 86, 148–156.

Safford, R.J., 2008. The dermal sensitisation threshold—A TTC approach for allergic contact dermatitis. *Regul. Toxicol. Pharmacol.* 51 (2), 195–200.

Safford, R.J., Api, A.M., Roberts, D.W., Lalko, J.F., 2015a. Extension of the dermal sensitization threshold (DST) approach to incorporate chemicals classified as reactive. *Regul. Toxicol. Pharmacol.* 72 (3), 694–701.

Safford, R.J., Aptula, A.O., Gilmour, N., 2011. Refinement of the dermal sensitisation threshold (DST) approach using a larger dataset and incorporating mechanistic chemistry domains. *Regul. Toxicol. Pharmacol.* 60 (2), 218–224.

Salvito, D.T., Senna, R.J., Federle, T.W., 2002. A Framework for prioritizing fragrance materials for aquatic risk assessment. *Environ. Toxicol. Chem.* 21 (6), 1301–1308.

Us Epa, 2012a. Estimation Programs Interface Suite for Microsoft Windows, v4.0–v4.11. United States Environmental Protection Agency, Washington, DC, USA.

Us Epa, 2012b. The ECOSAR (ECological Structure Activity Relationship) Class Program for Microsoft Windows, v1.11. United States Environmental Protection Agency, Washington, DC, USA.